


Neuron

Merkel Cells Activate Sensory Neural Pathways through Adrenergic Synapses

Graphical Abstract



Authors

Benjamin U. Hoffman,
Yoshichika Baba,
Theanne N. Griffith, ...,
Ardem Patapoutian, David Sulzer,
Ellen A. Lumpkin

Correspondence

ellen.a.lumpkin@gmail.com

In Brief

Hoffman et al. reveal the molecular machinery underlying neurotransmission at a gentle-touch receptor. Employing Eccles's classical criteria for a chemical synapse, they show that epithelial Merkel cells communicate with sensory neurons through β₂-adrenergic receptors at excitatory synapses.

Highlights

- Epidermal Merkel cells form adrenergic synapses with Aβ mechanosensory afferents
- Norepinephrine directly activates action potentials in Merkel-cell afferents
- Merkel cells employ SNARE-dependent vesicular release to excite tactile afferents
- Neuronal β₂-adrenergic receptors are required for slowly adapting type I responses



Merkel Cells Activate Sensory Neural Pathways through Adrenergic Synapses

Benjamin U. Hoffman,^{1,2} Yoshichika Baba,¹ Theanne N. Griffith,¹ Eugene V. Mosharov,³ Seung-Hyun Woo,^{4,8} Daniel D. Roybal,⁵ Gerard Karsenty,⁶ Ardem Patapoutian,⁴ David Sulzer,³ and Ellen A. Lumpkin^{1,2,7,9,*}

¹Department of Physiology & Cellular Biophysics, Columbia University, New York, NY, USA

²Program in Neurobiology & Behavior, Columbia University, New York, NY, USA

³Departments of Psychiatry, Neurology, and Pharmacology, Columbia University; Division of Molecular Therapeutics, New York State Psychiatric Institute, New York, NY, USA

⁴The Scripps Research Institute & Howard Hughes Medical Institute, La Jolla, CA, USA

⁵Pharmacology Graduate Program, Columbia University, New York, NY, USA

⁶Department of Genetics and Development, Columbia University, New York, NY, USA

⁷Department of Dermatology, Columbia University, New York, NY, USA

⁸Present address: Genomics Institute of the Novartis Research Foundation, San Diego, CA, USA

⁹Lead Contact

*Correspondence: ellen.a.lumpkin@gmail.com

<https://doi.org/10.1016/j.neuron.2018.10.034>

SUMMARY

Epithelial-neuronal signaling is essential for sensory encoding in touch, itch, and nociception; however, little is known about the release mechanisms and neurotransmitter receptors through which skin cells govern neuronal excitability. Merkel cells are mechanosensory epidermal cells that have long been proposed to activate neuronal afferents through chemical synaptic transmission. We employed a set of classical criteria for chemical neurotransmission as a framework to test this hypothesis. RNA sequencing of adult mouse Merkel cells demonstrated that they express presynaptic molecules and biosynthetic machinery for adrenergic transmission. Moreover, live-cell imaging directly demonstrated that Merkel cells mediate activity- and VMAT-dependent release of fluorescent catecholamine neurotransmitter analogs. Touch-evoked firing in Merkel-cell afferents was inhibited either by pre-synaptic silencing of SNARE-mediated vesicle release from Merkel cells or by neuronal deletion of β_2 -adrenergic receptors. Together, these results identify both pre- and postsynaptic mechanisms through which Merkel cells excite mechanosensory afferents to encode gentle touch.

INTRODUCTION


As a sensory-neural organ, skin provides both a protective barrier and an environmental interface that allows organisms to react to changing conditions. Signaling between epithelial cells and somatosensory neurons shape touch, itch, nociception, and chemoreception (Wilson et al., 2013; Maksimovic et al., 2014; Baumbauer et al., 2015; Pang et al., 2015; Moehring

et al., 2018; Bellono et al., 2017). Little is known, however, about the mechanisms through which skin cells release neuroactive molecules to govern neuronal excitability.

Since their initial description as “touch cells,” Merkel cells have served as the archetypical skin cell that mediates somatosensation (Merkel, 1875). These epithelial-derived cells complex with A β low-threshold mechanoreceptors (LTMRs) to produce slowly adapting type I (SAI) responses (Iggó and Muir, 1969; Woodbury and Koerber, 2007; Maricich et al., 2009; Morrison et al., 2009; Van Keymeulen et al., 2009). Recent work has demonstrated that Merkel cells mediate sustained SAI responses through Piezo2-dependent ion channels (Ikeda et al., 2014; Maksimovic et al., 2014; Woo et al., 2014). Moreover, optogenetics revealed that Merkel cells are necessary and sufficient to evoke sustained firing in A β LTMRs (Maksimovic et al., 2014). These studies establish Merkel cells as mechanosensory receptor cells, but the mechanisms through which these epidermal cells activate sensory neurons is still debated.

A long-standing model is that Merkel cells form chemical synapses with sensory afferents (Tachibana and Nawa, 2002; Maksimovic et al., 2013). Consistent with this hypothesis, Merkel cells are enriched in molecules that mediate synaptic vesicle release and are immunoreactive for neurotransmitters, neuromodulators, and neurotransmission machinery (Fantini and Johansson, 1995; García-Caballero et al., 1989; Hartschuh and Weihe, 1988; Weihe et al., 1998; Leung and Wong, 2000; Haerberle et al., 2004; Tachibana and Nawa, 2005; Maksimovic et al., 2013). At the ultrastructural level, Merkel cells form synaptic-like contacts with sensory terminals; however, these are marked by dense-core vesicles rather than clear-core vesicles that typically mediate fast synaptic transmission (Andres, 1966; Breathnach and Robins, 1970; Chen et al., 1973; Mihara et al., 1979). Several studies have proposed the Merkel cell-neurite complex is either serotonergic or glutamatergic (Fagan and Cahusac, 2001; He et al., 2003; Hitchcock et al., 2004; Cahusac et al., 2005; Press et al., 2010; Chang et al., 2016). Nonetheless, direct functional evidence is lacking to identify the Merkel cell's presynaptic mechanisms and the neurotransmitter receptor(s)





(legend on next page)

that act cell autonomously in sensory neurons. Thus, we sought to systematically dissect molecular mechanisms that mediate neurotransmission at the Merkel cell-neurite complex.

As a framework for this analysis, we turned to the work of Eccles, who proposed a set of fundamental criteria for a bona fide chemical synapse based on Dale and Loewi's pioneering studies (Dale, 1937; Loewi, 1937; Eccles, 1964): (1) biosynthetic and degradative mechanisms for chemical transmission must be present in the presynaptic cell, (2) neurotransmitter must be present in the presynaptic cell, (3) neurotransmitter must be released when the presynaptic cell is stimulated, (4) the postsynaptic cell must be activated (or inactivated) by direct application of the neurotransmitter, and (5) pharmacological antagonism of the neurotransmitter must block the physiological action of the postsynaptic cell. By evaluating these classical criteria, we demonstrate that Merkel cells form SNARE-dependent chemical synapses that excite touch-sensitive neurons through adrenergic receptors.

RESULTS

Merkel Cells Are Presynaptic, Catecholaminergic Cells

A handful of presynaptic proteins have been localized to Merkel cells; however, a genome-wide analysis of presynaptic signaling molecules in adult Merkel cells is lacking. We performed RNA sequencing (RNA-seq) on Merkel cells and keratinocytes purified from adult mice expressing an ATOH1-GFP fusion protein (*Atoh1^{GFP}*), which selectively marks Merkel cells in skin (Figure 1A; Haeberle et al., 2004; Rose et al., 2009; Woo et al., 2014). RNA-seq analysis identified 3,364 genes that were differentially expressed between Merkel cells and basal keratinocytes (Figures 1B and 1C; Table S1). Next, we generated a list of 542 genes encoding presynaptic molecules based on proteomic analyses of presynapses and gene ontology annotations (Table S2; Abul-Husn et al., 2009; Boyken et al., 2013; Aken et al., 2017; Blake et al., 2017). Of these, 123 genes were enriched in adult Merkel cells (Figure 1D), in agreement with previous studies of neonatal Merkel cells (Haerberle et al., 2004). Merkel-cell-enriched molecules span a wide range of presynaptic structures and functions, including the active zone, adhesion and cell surface, dense-core and synaptic vesicles, ion channels, transporters and receptors, neurotransmitter synthesis, and SNAREs. Thus, adult Merkel cells encode all the presynaptic machinery necessary for regulated release of neurotransmitters.

Dense-core vesicles package neuropeptides and small-molecule neurotransmitters, such as monoamines, which are capable of exciting action potentials in neurons (Araneda and Firestein,

2006; Ramírez-Franco et al., 2016; Zhou et al., 2016). Monoamines include tryptophan-derived serotonin (5HT) and tyrosine-derived catecholamines, such as dopamine, epinephrine, and norepinephrine. We noted that Merkel cells express *tyrosine hydroxylase* (*Th*), which encodes the rate-limiting enzyme in catecholamine synthesis (Figure 1D; Molinoff and Axelrod, 1971). To verify that Merkel cells express *Th*, we performed immunohistochemistry on full-thickness skin specimens from *Th^{GFP}* reporter mice (Figure 1E; Matsushita et al., 2002). Merkel cells and their A β LTMR afferents were identified by immunoreactivity to keratin 8 (K8) and neurofilament heavy chain (NFH), respectively (Vielkind et al., 1995; Maricich et al., 2009). Ninety-four percent of K8-positive Merkel cells expressed Th-GFP ($n = 406$ Merkel cells from two mice). Merkel cells were also immunoreactive for TH protein (Figure 1F). Along with *Th*, adult mouse Merkel cells express *Slc18a2* transcripts encoding VMAT2 (Table S2; 2.6 ± 0.1 ; RPKM, mean \pm SEM, $n = 2$ replicates), which preferentially loads catecholamines into synaptic vesicles, and *Maoa* transcripts (18.3 ± 1.3), which encodes a key neuronal enzyme for monoamine degradation (Erickson et al., 1996; Shih et al., 1999). Thus, Merkel cells possess the machinery to synthesize catecholamines and load them into secretory vesicles.

Which catecholamines might Merkel cells synthesize? We performed high-performance liquid chromatography (HPLC) coupled with electrochemical detection of monoamine neurotransmitters on whisker follicles, which are highly enriched in Merkel cells (Feigin et al., 2001; Larsen et al., 2002). Follicles were micro-dissected to reduce vascular contributions. To quantify the Merkel-cell-dependent neurotransmitter content, we compared whisker follicles from adult *K14^{Cre};Atoh1^{LacZ/ff}* mice, which lack Merkel cells, to littermate controls and found that norepinephrine was reduced by one-third in samples lacking Merkel cells (Figures S1A–S1C). Given that whisker follicles are highly vascularized, sympathetic innervation is a likely source of residual norepinephrine in mice lacking Merkel cells. Epinephrine and 5HT were comparable between genotypes, and dopamine was not detected. The lack of dopamine suggests that dopamine is efficiently converted to norepinephrine in the follicle. Indeed, Merkel cells were immunoreactive for dopamine beta hydroxylase (DBH), which synthesizes norepinephrine from dopamine (Figure S1D). These data suggest that epidermal Merkel cells are a significant source of norepinephrine, but not 5HT, epinephrine, or dopamine.

We were surprised to find that Merkel cells did not contribute to 5HT levels in whisker follicles, given that previous studies proposed that Merkel cells are serotonergic (He et al., 2003;

Figure 1. Merkel Cells Are Presynaptic, Catecholaminergic Cells

- (A) Epidermal cell purification strategy ($n = 2$ FACS purifications from $n \geq 2$, 7- to 8-week *Atoh1^{GFP}* mice each). FACS, fluorescence-activated cell sorting.
 (B) Volcano plot of genes differentially expressed between Merkel cells and keratinocytes. Dashed lines indicate $\log_2(\text{fold difference}) \geq 2$ and $P_{\text{adj}} < 0.01$ for differential expression (red, above threshold; black, below threshold).
 (C) Hierarchical clustering of differentially expressed genes. Rows represent RNA-seq replicates (R1/R2). Dendrograms show expression profiles of genes (top) and replicates (left). RPKM, reads per kilobase of exon per million reads mapped. Genes with RPKM < 2 across all samples are not displayed.
 (D) Presynaptic genes enriched in Merkel cells are grouped according to functional class. \log_2 -transformed fold difference \pm SE is plotted.
 (E) Axial projection of a whole-mount touch dome from an adult *Th^{GFP}* mouse stained with antibodies against NFH (blue in merge), K8 (magenta), and GFP (green).
 (F) Maximum projection of a touch dome in an epidermal peel stained with antibodies against K8 (magenta) and TH (green).
 Scale bars, 25 μm . See also Figure S1 and Tables S1 and S2.

Press et al., 2010; Chang et al., 2016, 2017). Consistent with our HPLC data, transcripts for the rate-limiting enzymes for 5HT biosynthesis were not detected in adult mouse Merkel cells (*Tph1*, 0.13 ± 0.02 ; *Tph2*, 0.01 ± 0.01 ; RPKM, mean \pm SEM). Moreover, analysis of a transgenic GFP reporter strain for the ionotropic 5HT_{3A} receptor revealed that NFH⁺ A β afferents that contacted mouse Merkel cells in touch domes and whisker follicles were not immunoreactive for 5HT_{3A}-GFP (Figure S1E; Gong et al., 2003). Instead, 5HT_{3A}-GFP-expressing afferents lacked NFH immunoreactivity, indicating that they are unmyelinated C fibers or thinly myelinated A δ fibers. These data are consistent with the role of 5HT₃ in nociception (Zeititz et al., 2002; Kayser et al., 2007). Thus, we conclude that Merkel cell-neurite complexes in adult mice do not form 5HT_{3A}-dependent serotonergic synapses.

In summary, these molecular and electrochemical results fulfill the first two of Eccles's criteria for a classical chemical synapse. Merkel cells express the biosynthetic and degradative machinery for adrenergic neurotransmitters. Moreover, the neurotransmitter norepinephrine in whisker follicles is dependent on Merkel cells. These results identify norepinephrine as a candidate neurotransmitter at Merkel cell-neurite complexes.

Merkel Cells Mediate Uptake and Touch-Stimulated Release of Fluorescent Neurotransmitters

We next asked whether Merkel cells package and release catecholamine neurotransmitters. We developed a semi-intact epidermal peel preparation that enables direct pharmacological and optical access to Merkel cells (Figure S2A). *Atoh1* reporter mice expressing GFP were used to identify Merkel cells, which formed readily visualized clusters in touch domes (Lumpkin et al., 2003; Maricich et al., 2009; Rose et al., 2009). We confirmed that Merkel cells are excitable in this preparation by depolarizing with high K⁺ Ringer's solution or by delivering focal displacements during calcium imaging experiments (Haeberle et al., 2004; Ikeda et al., 2014; Maksimovic et al., 2014). Both stimuli elicited rapid, reversible, and repeatable increases in cytoplasmic calcium (Figures S2B–S2I; Videos S1 and S2). Such calcium transients were abolished by antagonists of N- and P/Q-type (10 μ M ω -conotoxin MVII-C) and L-type (10 μ M nimodipine) voltage-gated Ca²⁺ channels (VGCCs; Figures S2J–S2M), in agreement with studies of dissociated Merkel cells (Haeberle et al., 2004, 2008).

To visualize neurotransmitter packaging, we employed a VMAT2-selective, neurotransmitter analog, fluorescent false neurotransmitter 206 (FFN206; Hu et al., 2013). Merkel cells reliably loaded FFN206 into puncta (328/330 Merkel cells from two animals) whereas surrounding keratinocytes showed no appreciable FFN206 accumulation. FFN206 puncta polarized to the cytoplasmic regions beneath nuclei (Figure 2A; Video S3), recapitulating the distribution of dense-core vesicles and presynaptic proteins near neuronal contacts (Chen et al., 1973; Hartschuh and Weihe, 1980; Haeberle et al., 2004). Reserpine, a selective VMAT2 antagonist, abolished FFN206 fluorescence in Merkel cells, demonstrating that VMAT2 activity is required for FFN206 loading (Figures 2B and 2C; Erickson et al., 1995). Thus, Merkel cells load VMAT2 substrates into subcellular puncta that likely represent secretory vesicle clusters.

To test whether Merkel cells are capable of evoked FFN206 release, we used live-cell imaging. Depolarization of FFN206-containing Merkel cells induced reliable destaining of FFN206 puncta (Figures 2D–2F), which was completely blocked by VGCC antagonists (Figures 2G–2I). Moreover, focal displacements caused rapid FFN206 destaining (Figure 2J–2L). Together, these data satisfy the functional release criterion of Eccles's classic chemical synapse: Merkel cells mediate evoked release of catecholaminergic vesicles in a VGCC-dependent manner.

Merkel Cells Employ SNARE-Dependent Vesicular Release to Mediate SAI Responses

We next used a synaptic silencing approach to directly test whether Merkel cells employ vesicle fusion for neurotransmission. Adult Merkel cells are highly enriched in vesicle and target SNARE transcripts, including vesicle-associated membrane protein 2 (*Vamp2*, 331 \pm 51; RPKM, Figure 1D). Tetanus neurotoxin light-chain subunit (TeNT) cleaves VAMP2 to block vesicle fusion at synapses (Yamamoto et al., 2003; Yu et al., 2004; Zhang et al., 2008; Chandrashekar et al., 2009). We crossed *K14^{Cre}* mice with *Rosa26^{flloxstop}TeNT-GFP* mice to selectively express TeNT in epidermal cells (*K14^{Cre};R26^{TeNT}*; Zhang et al., 2008; Morrison et al., 2009; Van Keymeulen et al., 2009). Control Merkel cells were enriched in VAMP2 protein compared with other epidermal cells (Figure 3A). TeNT reduced the proportion of VAMP2-positive Merkel cells by 3-fold (Figures 3A and 3B). Touch domes from *K14^{Cre};R26^{TeNT}* mice showed innervated Merkel-cell clusters, suggesting that vesicular neurotransmitter release is not needed for the development and maintenance of Merkel-cell innervation (Figure S3).

The functional consequences of epidermal-specific synaptic silencing were assessed with genotype-blind *ex vivo* single-unit recordings from Merkel-cell afferents, which were identified as fluorescently labeled, A β LTMR afferents that responded selectively to touch-dome indentation (Figures S4A–S4C, Figure 3C; Wellnitz et al., 2010). Littermate controls showed canonical SAI responses, characterized by high-frequency firing during dynamic stimulation and sustained low-frequency firing during static stimulation (Figure 3D). By contrast, Merkel-cell afferents from *K14^{Cre};R26^{TeNT}* mice (Figure 3D) produced responses with a 2-fold reduction in dynamic firing, a 3-fold reduction in static firing rates, and intermediately adapting (IA) firing patterns (Figures 3E and 3F). These responses phenocopied those from *Atoh1* conditional knockout mice, which completely lack Merkel cells, and epidermal-specific *Piezo2* knockout mice, which lack the Merkel cell's mechanotransduction channel (Maksimovic et al., 2014; Woo et al., 2014). Thus, SNARE-mediated vesicle release is required for the Merkel cell's contribution to SAI firing.

Norepinephrine Evokes Action Potentials in Merkel-Cell Afferents

Eccles's fourth criterion is that the postsynaptic cell must be activated by direct application of the neurotransmitter. Norepinephrine, dopamine, or 5HT was applied to receptive fields of Merkel-cell afferents during *ex vivo* skin-saphenous nerve recordings. Consistent with HPLC results, local application of




Figure 2. Merkel Cells Mediate Uptake and Activity-Evoked Release of Fluorescent Neurotransmitters

(A) Left: Merkel cells (magenta) loaded with FFN206 (1 μ M, cyan) in an epidermal peel preparation from an *Atoh1^{nGFP}* mouse, which expresses a nuclear localized GFP driven by *Atoh1* enhancer elements. Right: three-dimensional reconstruction, rotated. Images are representative of $n = 9$ touch domes, 2 animals.

(B) FFN206 loading in *Atoh1^{nGFP}* Merkel cells co-incubated with vehicle (left) or 1 μ M reserpine (right).

(C) Quantification of FFN206 loading (background subtracted fluorescence, one-way ANOVA, Tukey's post hoc; $^{\dagger}p < 0.0001$ for all comparisons; mean \pm SEM; $n = 205$ –331 Merkel cells per group, 2 animals).

(D–L) FFN206-loaded Merkel cells stimulated with high K^+ Ringer's solution. Pseudo-color images show the ratio of FFN206 fluorescence to baseline 240 s after application of high K^+ Ringer's solution. Scale indicates percent decrease in pixel intensity from baseline. (D) *Atoh1^{nGFP}*-positive Merkel cells (magenta). (E) Baseline normalized, intensity versus time traces of puncta indicated by arrowheads in (D). Black bar, high K^+ application. (F) Mean (red) \pm SEM (gray) of puncta that showed release in response to depolarization ($n = 40$ cells, 3 animals).

(G–I) Merkel cells loaded with FFN206, incubated with VGCC antagonists (10 μ M nimodipine + 10 μ M ω -conotoxin MVII-C), and stimulated with high K^+ Ringer's solution. (G) *Atoh1^{nGFP}*-positive Merkel cells (magenta) and pseudo-color difference image. (H) Baseline normalized, intensity versus time traces of puncta indicated by arrowheads in (G). Black bar, high K^+ application. (I) Mean (red) \pm SEM (gray) of summary data ($n = 69$ Merkel cells, 3 mice).

(J–L) Merkel cells loaded with FFN206 and stimulated with displacement. (J) Left: Merkel cells (*Atoh1^{nGFP}*, magenta). Right: ratio of FFN206 fluorescence to baseline 40 s after the onset of displacement. Dashed lines outline the stimulus probe. (K) Baseline normalized, intensity versus time traces of puncta indicated by the arrowhead in (J). Black bar, mechanical stimulation. (L) Mean (red) \pm SEM (gray) of mechanically evoked FFN206 release ($n = 6$ Merkel cells, 3 animals).

Scale bars, 20 μ m. See also [Figure S2](#) and [Videos S1, S2, and S3](#).

norepinephrine evokes firing in Merkel-cell afferents in the absence of touch stimuli ([Figures 4A–4C](#)). By contrast, neither dopamine nor 5HT activated action potentials in these afferents. Moreover, norepinephrine did not activate firing in A-fiber rapidly adapting (RA) LTMRs ([Figures S5A and S5B](#)). Thus, the excit-

atory effect of norepinephrine is not a general property of A-fiber LTMRs. Building on our Merkel-cell transcriptome analysis, which shows that Merkel cells are equipped to synthesize norepinephrine, these results directly demonstrate that norepinephrine preferentially excites Merkel-cell afferents.




Figure 3. Merkel Cells Employ SNARE-Dependent Vesicular Release to Mediate SAI Responses

(A) Images of Merkel cells (magenta) and VAMP2 (cyan) immunoreactivity from littermate control (R26^{TeNT}) and TeNT-expressing mice (K14^{Cre};R26^{TeNT}; scale bar, 10 μ m).

(B) Quantification of Merkel cells with detectable VAMP2 immunoreactivity (cyan; two-sided Fisher's exact test, $^{\dagger}p < 0.0001$; $n = 149$ –161 cells from 3 mice per group).

(C) Schematic of *ex vivo* skin-nerve recording preparation (saphenous nerve, red; FM1-43-labeled touch dome, green). Receptive fields of touch-dome afferents were identified by FM1-43 fluorescence in epidermal-side-up recordings.

(D) Recordings from touch-dome afferents from littermate-control (left) and K14^{Cre};R26^{TeNT} (right) mice. Top traces: displacement (dashed lines, point of skin contact). Middle traces: action potential trains. Bottom: instantaneous firing frequency (IFF) plots (blue region, dynamic [ramp] phase; gray region, static [late hold] phase).

(E–G). Maximal touch-evoked responses from littermate-control (cyan) and K14^{Cre};R26^{TeNT} (magenta). Peak dynamic firing rate (E) and mean static firing rate (F). Mean \pm SEM; two-tailed Mann-Whitney test, $^{\Omega}p < 0.005$. (G) Units were classified as slowly adapting (SA, white) if spikes were observed throughout the 5-s hold phase in $\geq 75\%$ of stimulus presentations; otherwise, they were classified as intermediately adapting (IA, black; two-sided Fisher's exact test, $^{\Omega}p < 0.005$; $n = 10$ –11 fibers, 9 mice per group). See also Figures S3 and S4.

Neuronal β_2 -adrenergic Receptors Mediate SAI Responses

Norepinephrine signals through metabotropic adrenergic receptors (ARs; Hein and Kobilka, 1995). Qualitative immunohistochemistry showed co-localization of NFH and β_2 AR immunoreactivity beneath some Merkel cells (Figure 5A). Moreover, single-molecule *in situ* hybridization revealed that a subset of dorsal root ganglia (DRG) neurons marked by *TrkC*^{tdTomato} and NFH, which includes Merkel-cell afferents, was enriched in β_2 AR transcripts (*Adrb2*; Figures S5C and S5D). Thus, β_2 ARs are well positioned to mediate signaling downstream of norepinephrine release from Merkel cells.

To test whether β_2 ARs are required for touch-evoked SAI responses, we locally applied the selective β_2 AR antagonist ICI 118,551 (henceforth referred to as ICI) to Merkel-cell afferents and delivered suprathreshold mechanical stimuli at 2-min intervals (Figure 5B). Compared with vehicle-treated controls, ICI suppressed dynamic SAI firing by 2-fold and static firing by 3.5-fold (Figures 5C–5F). The effects of ICI on static SAI firing partially reversed within 60 min of ICI washout (Figures S5E and S5F). Such incomplete washout from intact skin is not surprising, given the hydrophobic nature of ICI. Thus, touch-evoked SAI responses require activation of β_2 ARs.

We then used a genetic approach to confirm that β_2 ARs function cell autonomously in neurons to mediate SAI responses. We made *Wnt1*^{Cre};*Adrb2*^{fl/fl} mice, which harbor a deletion of β_2 ARs in neural crest derivatives, including somatosensory neurons (Jackson et al., 2012; Lewis et al., 2013). Touch domes from *Wnt1*^{Cre};*Adrb2*^{fl/fl} and littermate control mice showed comparable Merkel cell-nerve complexes, indicating that touch-dome innervation is normal in neuronal β_2 AR knockout mice (Figures S5G and S5H). Compared with typical SAI responses in littermate controls, Merkel-cell afferents from *Wnt1*^{Cre};*Adrb2*^{fl/fl} mice showed dramatically attenuated dynamic and static firing rates and IA responses (Figures 5G–5J). Indeed, responses from *Wnt1*^{Cre};*Adrb2*^{fl/fl} afferents were indistinguishable from those of *Atoh1* knockout mice lacking Merkel cells, mice whose Merkel cells lacked *Piezo2*, and TeNT mice with synaptically silenced Merkel cells (Figure 2; Figure S6). Together, these data demonstrate that neuronal β_2 ARs are required for touch-evoked SAI responses (i.e., the physiological action of the post-synaptic cell), satisfying Eccles's fifth criterion.

DISCUSSION

Epithelial-neuronal crosstalk plays an essential role in sensory signaling; however, our understanding of release mechanisms




Figure 4. Norepinephrine Excites Action Potentials in Merkel-Cell Afferents

(A–C) Norepinephrine (NE), serotonin (5HT), or dopamine (DA) was applied to receptive fields of touch-dome afferents in *ex vivo* skin-nerve recordings. Receptive fields of touch-dome afferents were identified by FM1-43 fluorescence and then preparations were flipped dermis-side up for receptive-field perfusion. (A) Left: neurotransmitter-evoked action potentials. Right: comparison of touch-evoked and neurotransmitter-evoked spike waveforms. Black bar indicates perfusion of 5 mM neurotransmitter (10 min). (B) Total spikes evoked by each neurotransmitter at 0.1 mM, 1.0 mM, or 5.0 mM

(two-way ANOVA with Tukey's post hoc, $^{\dagger}p < 0.0001$; $n_{NE} = 8$ units, 8 mice; $n_{5HT} = 9$ units, 9 mice; $n_{DA} = 4$ units, 4 mice; lines, medians). (C) Agonist-response relationship of NE-evoked spikes in Merkel-cell afferents (mean \pm SEM; EC₅₀ = 2.5 mM; $R^2 = 0.89$; $n = 11$ units, 9 mice).

and neurotransmitter receptors through which skin cells govern neuronal excitability is in its infancy. Adopting Eccles's definition of a bona fide chemical synapse, we demonstrate that epidermal Merkel cells excite the nervous system through SNARE-dependent adrenergic synapses. Our results show that Merkel cells express the biosynthetic and degradative machinery for adrenergic neurotransmission (criterion 1) and that norepinephrine in Merkel cell-rich skin areas depends on the presence of Merkel cells (criterion 2). In epidermis, Merkel cells selectively mediate VMAT- and VGCC-dependent evoked release of fluorescent catecholamines (criterion 3). Moreover, by using a tissue-specific genetic strategy to cleave Merkel-cell VAMP2, we directly demonstrate that SNARE-mediated vesicle release is essential for touch-evoked SAI responses. Turning to the postsynaptic cell, exogenous norepinephrine directly excites action potentials in Merkel-cell afferents (criterion 4), and touch-stimulated SAI responses are disrupted by pharmacological blockade or neuron-specific deletion of β_2 ARs (criterion 5). Together, these results identify both pre- and postsynaptic mechanisms through which touch-dome Merkel cells excite mechanosensory afferents (Figure 6).

The Merkel Cell-Neurite Complex Is an Adrenergic Synapse

Merkel cells are epithelial-derived mechanosensory cells whose role in gentle touch has recently been clarified. Mice harboring an epidermal-specific deletion of the neuronal transcription factor *Atoh1* lack Merkel cells throughout the body (Maricich et al., 2009; Morrison et al., 2009; Van Keymeulen et al., 2009). Electrophysiological analysis of these mice showed that Merkel cells potentiate SAI firing rates during dynamic and static touch and enable sustained firing during static pressure (Maksimovic et al., 2014). Moreover, sustained firing requires the mechanically gated ion channel Piezo2 in Merkel cells (Figure S6; Ikeda et al., 2014; Maksimovic et al., 2014; Woo et al., 2014). Thus, Merkel cells enhance the excitability of A β LTMRs during dynamic touch stimuli and mediate mechanosensory responses to gentle pressure.

Merkel cells accomplish these functions by signaling to sensory neurons at adrenergic synapses. By silencing SNARE-mediated vesicle release in epidermis, we found that the responses of Merkel-cell afferents recapitulate the suppressed dynamic firing and truncated static firing observed in

epidermal-specific *Atoh1* and *Piezo2* knockout mice (Maksimovic et al., 2014; Woo et al., 2014). These data are consistent with a recent report that bath-applied Botulinum toxin reduced compound action potentials in whisker afferents (Chang et al., 2016). Given that Merkel cells are highly enriched in VAMP2, the effects of epidermal TeNT expression are most likely due to VAMP2 cleavage in Merkel cells. Likewise, neuronal deletion of β_2 ARs phenocopied the effects of Merkel-cell synaptic silencing. Together, these results suggest that SNARE-dependent neurotransmitter release followed by β_2 AR activation on neurons accounts for the Merkel cell's role in SAI firing (Figure 6).

This finding is surprising, given the plethora of candidate neuromodulators at the Merkel cell-neurite complex (Maksimovic et al., 2013). Neuropeptides such as CCK, VIP, Met-Enkephalin, and substance P are present in Merkel cells (Hartschuh et al., 1979; Fantini and Johansson, 1995; Haeberle et al., 2004), and our transcriptome analysis demonstrates that Merkel cells are fully equipped to release these peptides through dense-core vesicles. Neurotransmitters such as ATP and glutamate have also been proposed to mediate Merkel-cell signaling (Toyoshima and Shimamura, 1991; Nakamura and Strittmatter, 1996; Fagan and Cahusac, 2001; Haeberle et al., 2004; Hitchcock et al., 2004; Nunzi et al., 2004). As previous functional studies have used pharmacology in semi-intact preparations, it is unclear whether these transmitters mediate autocrine signaling in Merkel cells or paracrine signaling to other cell types in skin. Consistent with autocrine signaling, glutamate and ATP receptors are expressed in Merkel cells (Table S1; Haeberle et al., 2004). By contrast, we observed β_2 AR immunoreactivity at the terminals of Merkel-cell afferents, and neuronal deletion of β_2 AR indicates that adrenergic signaling acts intrinsically in neurons to enable SAI responses. Intriguingly, neuroanatomical studies suggest that Merkel cells are contacted by neurons other than A β SAI afferents, including thinly myelinated and unmyelinated fibers (Reinisch and Tschachler, 2005; Lesniak et al., 2014; Niu et al., 2014). Whether Merkel cells communicate with these fiber types through β_2 ARs or other signaling pathways is unknown.

5HT, which has well-established roles in itch and nociception, has also been investigated in Merkel cells. Amperometry detected mechanically stimulated monoamine release from rodent whisker follicles (Chang et al., 2016); however, the methods employed do not distinguish 5HT from catecholamines (Pennington




Figure 5. Neuronal β_2 ARs Are Required for Touch-Evoked SAI Responses

(A) Merkel cells (K8, magenta), Merkel-cell afferent (NFH, cyan), and β_2 AR (yellow) immunoreactivity in adult mouse back skin (scale bar, 25 μ m).

(B) Recordings from Merkel-cell afferents and corresponding IFF plots before and 30 min after treatment with vehicle or 50 μ M ICI 118,551. Traces and shading as in Figure 3D.

(C–F) Peak dynamic firing rate (C and D) and mean static firing rates (E and F) from afferents treated with vehicle (cyan) or 50 μ M ICI 118,551 (magenta) and stimulated at 2-min intervals. (C and E) Firing rates at each time point were normalized to $t = 0$ to average responses across units (mean \pm SEM). Black bar indicates time of vehicle or 50 μ M ICI 118,551 application. (D and F) Firing rates of all units before ($t = 0$) and after ($t = 30$) treatment with vehicle or 50 μ M ICI 118,551. Two-way matched ANOVA with Sidak's post hoc; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, † $p < 0.0001$; $n = 5$ –6 units from 5–6 mice per group.

(G) Recordings from Merkel-cell afferents from littermate control (left) and *Wnt1^{Cre};Adrb2^{fl/fl}* (right) mice.

(H–J) Maximal touch-evoked responses from littermate control (cyan) and *Wnt1^{Cre};Adrb2^{fl/fl}* (magenta) units. (H) Peak dynamic firing rate. (I) Mean static firing rate (two-tailed Student's t test, *** $p < 0.001$; mean \pm SEM). (J) Proportion of Merkel-cell afferents with SA (white) versus IA firing patterns (black); two-sided Fisher's exact test, † $p < 0.005$; $n = 9$ –10 units, 9 mice per group). See also Figure S5.

et al., 2004). Moreover, transcriptome analysis in this study indicates that adult mouse Merkel cells express the rate-limiting enzyme for catecholamine synthesis but lack enzymes that produce 5HT. Moreover, 5HT did not excite action potentials in Merkel-cell afferents innervating touch domes, whereas norepinephrine elicited robust firing. These data stand in contrast with recent studies of 5HT signaling in mouse whisker afferents, which used suction electrode recordings of nerve bundles without spike sorting to isolate single-unit responses (Chang et al., 2016, 2017). Given that 5HT receptors are broadly expressed among sensory neurons (Zeitz et al., 2002; Lin et al., 2011; Urtikova et al., 2012; Morita et al., 2015; Salzer et al., 2016), it is likely that many types of somatosensory neurons contribute to 5HT-evoked compound action potentials. Consistent with this interpretation, mouse whisker follicles receive

our data do not support an excitatory role for 5HT signaling in mouse touch-dome afferents, Merkel cells might produce 5HT in other species or skin structures. An intriguing possibility is that Merkel-cell-derived 5HT could potentiate NFH-negative sensory neurons in pathological states such as chronic pain and itch. Indeed, a role for Merkel cells in mechanically evoked itch has been recently proposed (Feng et al., 2018).

Adrenergic Signaling in the Somatosensory System

Previous studies of adrenergic signaling in the somatosensory system have focused on nociception. In the descending antinociceptive system, for example, noradrenergic neurons in the locus coeruleus project to the dorsal horn of the spinal cord to inhibit nociceptive signals (Westlund and Coulter, 1980; Clark and Proudfoot, 1991; Fields et al., 1991). In the periphery, the role of




Figure 6. Model for Adrenergic Synaptic Transmission at the Merkel Cell-Neurite Complex

Touch activates cation influx through Piezo2 channels to depolarize Merkel cells and their afferents. During dynamic stimuli, neuronal Piezo2 stimulates action potential firing. Merkel-cell depolarization results in voltage-dependent calcium channel activation and SNARE-mediated release of norepinephrine. Norepinephrine binds to β_2 -adrenergic receptors on SAI afferents, which increases dynamic firing rates and initiates action potentials during static stimulation. See also Figure S6.

adrenergic signaling in nociception has been attributed to the sympathetic nervous system. After peripheral nerve injury or inflammation, sympathetic neurons sprout within DRG to sensitize nociceptive neurons to adrenergic stimulation (Sato and Perl, 1991; McLachlan et al., 1993; Bossut et al., 1996; Leem et al., 1997). Moreover, adrenergic compounds induce nociceptive responses in injured or inflamed skin (Chabal et al., 1992; Torebjörk et al., 1995; Choi and Rowbotham, 1997; Khasar et al., 1999). By contrast, norepinephrine does not elicit mechanonociceptive responses in naive tissue (Fuchs et al., 2001). To our knowledge, adrenergic signaling has not been implicated in the encoding of gentle touch. Thus, our results define a previously unsuspected paradigm for adrenergic signaling in somatosensation.

β_2 ARs and Neuronal Excitability

How might norepinephrine signaling through metabotropic β_2 ARs excite action potentials in postsynaptic A β LTMRs? A pre-

cedent for an excitatory metabotropic synapse is set by the retinal photoreceptor-ON bipolar cell synapse, which excites postsynaptic neurons via mGluR6 receptors coupled through G_{α_o} signaling to TRPM1-dependent cation channels (Martemyanov and Sampath, 2017).

β_2 ARs couple to G_{α_s} signaling, which controls neuronal excitability through altering the activity of ion channels. G_{α_s} activation stimulates adenylyl cyclase to increase adenosine 3',5'-cyclic monophosphate (cAMP) and activate downstream effectors such as cAMP-dependent protein kinase A (PKA; Rosenbaum et al., 2009), both of which can modulate ion channels. For example, in olfactory sensory neurons, a $G_{\alpha_{olf/s}}$ -coupled sensory transduction cascade culminates in gating of cyclic nucleotide-gated (CNG) cation channels (Nakamura and Gold, 1987). CNG channels likewise function as a critical component of membrane depolarization in vertebrate phototransduction (Yau and Baylor, 1989) and underlie inward conductances in hippocampal neurons (Leinders-Zufall et al., 1995). Intriguingly, CNG channel expression is detected in NFH-positive DRG neurons (Usoskin et al., 2015). Intracellular cAMP also signals through hyperpolarization-activated cyclic nucleotide-gated (HCN) cation channels (Mayer and Westbrook, 1983). Currents mediated by HCN channels are found in many types of sensory neurons, including A β LTMRs (Gao et al., 2012; Chang et al., 2017). Finally, β_2 ARs form a protein complex with $Ca_v1.2$ channels, which provides spatiotemporal coupling of β_2 ARs to calcium channel activation (Davare et al., 2001). PKA-mediated phosphorylation of $Ca_v1.2$ contributes to an increase in inward Ca^{2+} conductance (Hell et al., 1993; Gao et al., 1997; Bünemann et al., 1999). Such cAMP-PKA-mediated modulation of ion channels can occur with a latency of 500 ms (Lancaster et al., 2006). Additionally, the compartmentalization of cAMP signals into microdomains by tonic phosphodiesterase activity has been proposed to enable local activation of cAMP-PKA signaling (Rich et al., 2000; Jurevicius et al., 2003). Together, these mechanisms suggest a plausible model of norepinephrine-evoked, β_2 AR-mediated activation of A β LTMRs with fast kinetics; however, future studies are needed to establish whether these signaling mechanisms mediate excitation in Merkel-cell afferents.

β_2 -adrenergic Signaling in Health and Disease

β_2 -adrenergic signaling is a cornerstone therapeutic target for the treatment of cardiovascular diseases. β -blockers, which target β -adrenergic receptors, have been among the most widely used medicines for over half a century. A side effect of β -blocker therapy is paresthesia, or numbness and tingling (Van Buskirk, 1980; Stewart and Castelli, 1996). These symptoms are attributed to the vascular effects of β -blockers in peripheral tissues; however, our findings suggest that paresthesia might be a result of β -blockers disrupting adrenergic signaling in sensory afferents.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING

- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Animals
- **METHOD DETAILS**
 - RNA Sequencing and Analysis
 - Semi-intact Epidermal Peel Preparation
 - Immunohistochemistry
 - *In Situ* Hybridization
 - High-Performance Liquid Chromatography (HPLC)
 - FM1-43 Injections
 - *Ex Vivo* Skin-Nerve Electrophysiology
 - Calcium Imaging
 - FFN Imaging
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
- **DATA AND SOFTWARE AVAILABILITY**

SUPPLEMENTAL INFORMATION

Supplemental Information includes six figures, three tables, and three movies and can be found with this article online at <https://doi.org/10.1016/j.neuron.2018.10.034>.

A video abstract is available at <https://doi.org/10.1016/j.neuron.2018.10.034#mmc8>.

ACKNOWLEDGMENTS

Thanks to Dr. Eric Larson, Dr. Sue Kinnamon, Dr. Tom Finger, and Dr. Charles Zuker for advice and reagents; Dr. Joriene de Nooij for sharing neuronal gene expression data prior to publication; Dr. Kara Marshall for helpful preliminary data; Mr. Adan Horta for advice on transcriptome analysis; and members of the Lumpkin laboratory for discussions and comments on the manuscript. This research was supported by NIAMS (R01AR051219, E.A.L.) and NINDS (F31NS105449, B.U.H.). Authors are funded by the NIH (R01NS095435, R01MH108186, and R01DA07418, D.S.), Burroughs Wellcome Fund PDEP program (T.N.G.), NIGMS (T32GM007367, B.U.H.), JPB Foundation (D.S.), and the Howard Hughes Medical Institute (A.P.). RNA-seq was performed with support from the Genomics Institute of the Novartis Research Foundation. Core facilities were supported by NIAMS P30AR069632 and NCI P30CA013696. Funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTIONS

Conceptualization, B.U.H. and E.A.L.; Methodology, B.U.H., Y.B., T.N.G., E.V.M., S.-H.W., A.P., D.S., and E.A.L.; Software, B.U.H. and Y.B.; Formal Analysis, B.U.H., T.N.G., and E.A.L.; Investigation, B.U.H., Y.B., T.N.G., E.V.M., S.-H.W., and D.D.R.; Resources, G.K.; Data Curation, B.U.H. and E.A.L.; Writing – Original Draft, B.U.H. and E.A.L.; Writing – Review & Editing, B.U.H., S.-H.W., T.N.G., E.V.M., G.K., A.P., D.S., and E.A.L.; Visualization, B.U.H. and T.N.G.; Supervision, E.A.L.; Project Administration, E.A.L.; Funding Acquisition, B.U.H., E.A.L., and A.P.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: May 10, 2018

Revised: September 21, 2018

Accepted: October 22, 2018

Published: November 8, 2018

REFERENCES

Abul-Husn, N.S., Bushlin, I., Morón, J.A., Jenkins, S.L., Dolios, G., Wang, R., Iyengar, R., Ma'ayan, A., and Devi, L.A. (2009). Systems approach to explore components and interactions in the presynapse. *Proteomics* 9, 3303–3315.

Aken, B.L., Achuthan, P., Akanni, W., Amode, M.R., Bernsdorff, F., Bhai, J., Billis, K., Carvalho-Silva, D., Cummins, C., Clapham, P., et al. (2017). Ensembl 2017. *Nucleic Acids Res.* 45 (D1), D635–D642.

Andres, K.H. (1966). [On the fine structure of Merkel's tactile apparatus in the sinus hair]. *Naturwissenschaften* 53, 706.

Araneda, R.C., and Firestein, S. (2006). Adrenergic enhancement of inhibitory transmission in the accessory olfactory bulb. *J. Neurosci.* 26, 3292–3298.

Bai, L., Lehnert, B.P., Liu, J., Neubarth, N.L., Dickendesher, T.L., Nwe, P.H., Cassidy, C., Woodbury, C.J., and Ginty, D.D. (2015). Genetic identification of an expansive mechanoreceptor sensitive to skin stroking. *Cell* 163, 1783–1795.

Basbaum, A.I., Bautista, D.M., Scherrer, G., and Julius, D. (2009). Cellular and molecular mechanisms of pain. *Cell* 139, 267–284.

Baumbauer, K.M., DeBerry, J.J., Adelman, P.C., Miller, R.H., Hachisuka, J., Lee, K.H., Ross, S.E., Koerber, H.R., Davis, B.M., and Albers, K.M. (2015). Keratinocytes can modulate and directly initiate nociceptive responses. *eLife* 4, e09674.

Bellono, N.W., Bayrer, J.R., Leitch, D.B., Castro, J., Zhang, C., O'Donnell, T.A., Brierley, S.M., Ingraham, H.A., and Julius, D. (2017). Enterochromaffin cells are gut chemosensors that couple to sensory neural pathways. *Cell* 170, 185–198.e16.

Ben-Arie, N., Hassan, B.A., Birmingham, N.A., Malicki, D.M., Armstrong, D., Matzuk, M., Bellen, H.J., and Zoghbi, H.Y. (2000). Functional conservation of atonal and Math1 in the CNS and PNS. *Development* 127, 1039–1048.

Blake, J.A., Eppig, J.T., Kadin, J.A., Richardson, J.E., Smith, C.L., and Bult, C.J.; the Mouse Genome Database Group (2017). Mouse Genome Database (MGD)-2017: community knowledge resource for the laboratory mouse. *Nucleic Acids Res.* 45 (D1), D723–D729.

Bossut, D.F., Shea, V.K., and Perl, E.R. (1996). Sympathectomy induces adrenergic excitability of cutaneous C-fiber nociceptors. *J. Neurophysiol.* 75, 514–517.

Boyken, J., Grønberg, M., Riedel, D., Urlaub, H., Jahn, R., and Chua, J.J. (2013). Molecular profiling of synaptic vesicle docking sites reveals novel proteins but few differences between glutamatergic and GABAergic synapses. *Neuron* 78, 285–297.

Breathnach, A.S., and Robins, J. (1970). Ultrastructural observations on Merkel cells in human foetal skin. *J. Anat.* 106, 411.

Bünemann, M., Gerhardstein, B.L., Gao, T., and Hosey, M.M. (1999). Functional regulation of L-type calcium channels via protein kinase A-mediated phosphorylation of the beta(2) subunit. *J. Biol. Chem.* 274, 33851–33854.

Cahusac, P.M., Senok, S.S., Hitchcock, I.S., Genever, P.G., and Baumann, K.I. (2005). Are unconventional NMDA receptors involved in slowly adapting type I mechanoreceptor responses? *Neuroscience* 133, 763–773.

Chabal, C., Jacobson, L., Russell, L.C., and Burchiel, K.J. (1992). Pain response to perineuronal injection of normal saline, epinephrine, and lidocaine in humans. *Pain* 49, 9–12.

Chandrashekar, J., Yarmolinsky, D., von Buchholtz, L., Oka, Y., Sly, W., Ryba, N.J., and Zuker, C.S. (2009). The taste of carbonation. *Science* 326, 443–445.

Chang, W., Kanda, H., Ikeda, R., Ling, J., DeBerry, J.J., and Gu, J.G. (2016). Merkel disc is a serotonergic synapse in the epidermis for transmitting tactile signals in mammals. *Proc. Natl. Acad. Sci. USA* 113, E5491–E5500.

Chang, W., Kanda, H., Ikeda, R., Ling, J., and Gu, J.G. (2017). Serotonergic transmission at Merkel discs: modulation by exogenously applied chemical messengers and involvement of I_h currents. *J. Neurochem.* 141, 565–576.

Chen, S.Y., Gerson, S., and Meyer, J. (1973). The fusion of Merkel cell granules with a synapse-like structure. *J. Invest. Dermatol.* 61, 290–292.

Choi, B., and Rowbotham, M.C. (1997). Effect of adrenergic receptor activation on post-herpetic neuralgia pain and sensory disturbances. *Pain* 69, 55–63.

Clark, F.M., and Proudfoot, H.K. (1991). The projection of locus coeruleus neurons to the spinal cord in the rat determined by anterograde tracing combined with immunocytochemistry. *Brain Res.* 538, 231–245.

- Dale, H. (1937). Some recent extensions of the chemical transmission of the effects of nerve impulses. In *Les Prix Nobel en 1936* (Imprimerie Royale P.A. Norstedt & Söner), pp. 1–12.
- Dassule, H.R., Lewis, P., Bei, M., Maas, R., and McMahon, A.P. (2000). Sonic hedgehog regulates growth and morphogenesis of the tooth. *Development* 127, 4775–4785.
- Davare, M.A., Avdonin, V., Hall, D.D., Peden, E.M., Burette, A., Weinberg, R.J., Horne, M.C., Hoshi, T., and Hell, J.W. (2001). A beta2 adrenergic receptor signaling complex assembled with the Ca²⁺ channel Cav1.2. *Science* 293, 98–101.
- Drew, L.J., Wood, J.N., and Cesare, P. (2002). Distinct mechanosensitive properties of capsaicin-sensitive and -insensitive sensory neurons. *J. Neurosci.* 22, RC228.
- Eccles, J.C. (1964). *The Physiology of Synapses* (Academic Press Inc., Publishers).
- Erickson, J.D., Eiden, L.E., Schafer, M.K., and Weihe, E. (1995). Reserpine- and tetrabenazine-sensitive transport of (3)H-histamine by the neuronal isoform of the vesicular monoamine transporter. *J. Mol. Neurosci.* 6, 277–287.
- Erickson, J.D., Schafer, M.K., Bonner, T.I., Eiden, L.E., and Weihe, E. (1996). Distinct pharmacological properties and distribution in neurons and endocrine cells of two isoforms of the human vesicular monoamine transporter. *Proc. Natl. Acad. Sci. USA* 93, 5166–5171.
- Fagan, B.M., and Cahusac, P.M. (2001). Evidence for glutamate receptor mediated transmission at mechanoreceptors in the skin. *Neuroreport* 12, 341–347.
- Fantini, F., and Johansson, O. (1995). Neurochemical markers in human cutaneous Merkel cells. An immunohistochemical investigation. *Exp. Dermatol.* 4, 365–371.
- Feigin, A., Fukuda, M., Dhawan, V., Przedborski, S., Jackson-Lewis, V., Mentis, M.J., Moeller, J.R., and Eidelberg, D. (2001). Metabolic correlates of levodopa response in Parkinson's disease. *Neurology* 57, 2083–2088.
- Feng, J., Luo, J., Yang, P., Du, J., Kim, B.S., and Hu, H. (2018). Piezo2 channel-Merkel cell signaling modulates the conversion of touch to itch. *Science* 360, 530–533.
- Fields, H.L., Heinricher, M.M., and Mason, P. (1991). Neurotransmitters in nociceptive modulatory circuits. *Annu. Rev. Neurosci.* 14, 219–245.
- Fuchs, P.N., Meyer, R.A., and Raja, S.N. (2001). Heat, but not mechanical hyperalgesia, following adrenergic injections in normal human skin. *Pain* 90, 15–23.
- Gao, T., Yatani, A., Dell'Acqua, M.L., Sako, H., Green, S.A., Dascal, N., Scott, J.D., and Hosey, M.M. (1997). cAMP-dependent regulation of cardiac L-type Ca²⁺ channels requires membrane targeting of PKA and phosphorylation of channel subunits. *Neuron* 19, 185–196.
- Gao, L.L., McMullan, S., Djouhri, L., Acosta, C., Harper, A.A., and Lawson, S.N. (2012). Expression and properties of hyperpolarization-activated current in rat dorsal root ganglion neurons with known sensory function. *J. Physiol.* 590, 4691–4705.
- García-Caballero, T., Gallego, R., Rosón, E., Basanta, D., Morel, G., and Beiras, A. (1989). Localization of serotonin-like immunoreactivity in the Merkel cells of pig snout skin. *Anat. Rec.* 225, 267–271.
- Gong, S., Zheng, C., Doughty, M.L., Losos, K., Didkovsky, N., Schambra, U.B., Nowak, N.J., Joyner, A., Leblanc, G., Hatten, M.E., and Heintz, N. (2003). A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. *Nature* 425, 917–925.
- Haeblerle, H., Fujiwara, M., Chuang, J., Medina, M.M., Panditrao, M.V., Bechstedt, S., Howard, J., and Lumpkin, E.A. (2004). Molecular profiling reveals synaptic release machinery in Merkel cells. *Proc. Natl. Acad. Sci. USA* 101, 14503–14508.
- Haeblerle, H., Bryan, L.A., Vadakkan, T.J., Dickinson, M.E., and Lumpkin, E.A. (2008). Swelling-activated Ca²⁺ channels trigger Ca²⁺ signals in Merkel cells. *PLoS ONE* 3, e1750.
- Hartschuh, W., and Weihe, E. (1980). Fine structural analysis of the synaptic junction of Merkel cell-axon-complexes. *J. Invest. Dermatol.* 75, 159–165.
- Hartschuh, W., and Weihe, E. (1988). Multiple messenger candidates and marker substance in the mammalian Merkel cell-axon complex: a light and electron microscopic immunohistochemical study. *Prog. Brain Res.* 74, 181–187.
- Hartschuh, W., Weihe, E., Büchler, M., Helmstaedter, V., Feurle, G.E., and Forssmann, W.G. (1979). Met-enkephalin-like immunoreactivity in Merkel cells. *Cell Tissue Res.* 201, 343–348.
- He, L., Tuckett, R.P., and English, K.B. (2003). 5-HT₂ and 3 receptor antagonists suppress the response of rat type I slowly adapting mechanoreceptor: an in vitro study. *Brain Res.* 969, 230–236.
- Hein, L., and Kobilka, B.K. (1995). Adrenergic receptor signal transduction and regulation. *Neuropharmacology* 34, 357–366.
- Hell, J.W., Yokoyama, C.T., Wong, S.T., Warner, C., Snutch, T.P., and Catterall, W.A. (1993). Differential phosphorylation of two size forms of the neuronal class C L-type calcium channel alpha 1 subunit. *J. Biol. Chem.* 268, 19451–19457.
- Hinoi, E., Gao, N., Jung, D.Y., Yadav, V., Yoshizawa, T., Myers, M.G., Jr., Chua, S.C., Jr., Kim, J.K., Kaestner, K.H., and Karsenty, G. (2008). The sympathetic tone mediates leptin's inhibition of insulin secretion by modulating osteocalcin bioactivity. *J. Cell Biol.* 183, 1235–1242.
- Hitchcock, I.S., Genever, P.G., and Cahusac, P.M. (2004). Essential components for a glutamatergic synapse between Merkel cell and nerve terminal in rats. *Neurosci. Lett.* 362, 196–199.
- Hoffman, B.U., and Lumpkin, E.A. (2018). *buh2003/SpikeSorting PCA_DBSCAN: SpikeSortingPCA_DBSCAN v1.0.1 (Version v1.0.01, Zenodo)*.
- Hosogi, M., Schmelz, M., Miyachi, Y., and Ikoma, A. (2006). Bradykinin is a potent pruritogen in atopic dermatitis: a switch from pain to itch. *Pain* 126, 16–23.
- Hu, G., Henke, A., Karpowicz, R.J., Jr., Sonders, M.S., Farrimond, F., Edwards, R., Sulzer, D., and Sames, D. (2013). New fluorescent substrate enables quantitative and high-throughput examination of vesicular monoamine transporter 2 (VMAT2). *ACS Chem. Biol.* 8, 1947–1954.
- Iggo, A., and Muir, A.R. (1969). The structure and function of a slowly adapting touch corpuscle in hairy skin. *J. Physiol.* 200, 763–796.
- Ikeda, R., Cha, M., Ling, J., Jia, Z., Coyle, D., and Gu, J.G. (2014). Merkel cells transduce and encode tactile stimuli to drive Aβ-afferent impulses. *Cell* 157, 664–675.
- Jackson, C.R., Ruan, G.X., Aseem, F., Abey, J., Gamble, K., Stanwood, G., Palmiter, R.D., Iuvone, P.M., and McMahon, D.G. (2012). Retinal dopamine mediates multiple dimensions of light-adapted vision. *J. Neurosci.* 32, 9359–9368.
- Julius, D., and Basbaum, A.I. (2001). Molecular mechanisms of nociception. *Nature* 413, 203–210.
- Jurevicius, J., Skeberdis, V.A., and Fischmeister, R. (2003). Role of cyclic nucleotide phosphodiesterase isoforms in cAMP compartmentation following beta2-adrenergic stimulation of ICa_L in frog ventricular myocytes. *J. Physiol.* 551, 239–252.
- Kayser, V., Elfassi, I.E., Aubel, B., Melfort, M., Julius, D., Gingrich, J.A., Hamon, M., and Bourgoin, S. (2007). Mechanical, thermal and formalin-induced nociception is differentially altered in 5-HT1A^{-/-}, 5-HT1B^{-/-}, 5-HT2A^{-/-}, 5-HT3A^{-/-} and 5-HTT^{-/-} knock-out male mice. *Pain* 130, 235–248.
- Khasar, S.G., McCarter, G., and Levine, J.D. (1999). Epinephrine produces a beta-adrenergic receptor-mediated mechanical hyperalgesia and in vitro sensitization of rat nociceptors. *J. Neurophysiol.* 81, 1104–1112.
- Lancaster, B., Hu, H., Gibb, B., and Storm, J.F. (2006). Kinetics of ion channel modulation by cAMP in rat hippocampal neurones. *J. Physiol.* 576, 403–417.
- Larsen, K.E., Fon, E.A., Hastings, T.G., Edwards, R.H., and Sulzer, D. (2002). Methamphetamine-induced degeneration of dopaminergic neurons involves autophagy and upregulation of dopamine synthesis. *J. Neurosci.* 22, 8951–8960.
- Leem, J.W., Gwak, Y.S., Nam, T.S., and Paik, K.S. (1997). Involvement of alpha2-adrenoceptors in mediating sympathetic excitation of injured dorsal

- root ganglion neurons in rats with spinal nerve ligation. *Neurosci. Lett.* 234, 39–42.
- Leinders-Zufall, T., Rosenboom, H., Barnstable, C.J., Shepherd, G.M., and Zufall, F. (1995). A calcium-permeable cGMP-activated cation conductance in hippocampal neurons. *Neuroreport* 6, 1761–1765.
- Lesniak, D.R., Marshall, K.L., Wellnitz, S.A., Jenkins, B.A., Baba, Y., Rasband, M.N., Gerling, G.J., and Lumpkin, E.A. (2014). Computation identifies structural features that govern neuronal firing properties in slowly adapting touch receptors. *eLife* 3, e01488.
- Leung, M.S., and Wong, C.C. (2000). Expressions of putative neurotransmitters and neuronal growth related genes in Merkel cell-neurite complexes of the rats. *Life Sci.* 66, 1481–1490.
- Lewis, A.E., Vasudevan, H.N., O’Neill, A.K., Soriano, P., and Bush, J.O. (2013). The widely used Wnt1-Cre transgene causes developmental phenotypes by ectopic activation of Wnt signaling. *Dev. Biol.* 379, 229–234.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R.; 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079.
- Lin, S.Y., Chang, W.J., Lin, C.S., Huang, C.Y., Wang, H.F., and Sun, W.H. (2011). Serotonin receptor 5-HT2B mediates serotonin-induced hyperalgesia. *J. Neurosci.* 31, 1410–1418.
- Loewi, O. (1937). Die chemische Übertragung der Nervenwirkung. In *Les Prix Nobel en 1936* (Imprimerie Royale P.A. Norstedt & Söner), pp. 1–14.
- Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.
- Lumpkin, E.A., Collisson, T., Parab, P., Omer-Abdalla, A., Haeberle, H., Chen, P., Doetzelhofer, A., White, P., Groves, A., Segil, N., and Johnson, J.E. (2003). Math1-driven GFP expression in the developing nervous system of transgenic mice. *Gene Expr. Patterns* 3, 389–395.
- Maksimovic, S., Baba, Y., and Lumpkin, E.A. (2013). Neurotransmitters and synaptic components in the Merkel cell-neurite complex, a gentle-touch receptor. *Ann. N Y Acad. Sci.* 1279, 13–21.
- Maksimovic, S., Nakatani, M., Baba, Y., Nelson, A.M., Marshall, K.L., Wellnitz, S.A., Firozi, P., Woo, S.H., Ranade, S., Patapoutian, A., and Lumpkin, E.A. (2014). Epidermal Merkel cells are mechanosensory cells that tune mammalian touch receptors. *Nature* 509, 617–621.
- Maricich, S.M., Wellnitz, S.A., Nelson, A.M., Lesniak, D.R., Gerling, G.J., Lumpkin, E.A., and Zoghbi, H.Y. (2009). Merkel cells are essential for light-touch responses. *Science* 324, 1580–1582.
- Martemyanov, K.A., and Sampath, A.P. (2017). The transduction cascade in retinal ON-bipolar cells: signal processing and disease. *Annu. Rev. Vis. Sci.* 3, 25–51.
- Matsushita, N., Okada, H., Yasoshima, Y., Takahashi, K., Kiuchi, K., and Kobayashi, K. (2002). Dynamics of tyrosine hydroxylase promoter activity during midbrain dopaminergic neuron development. *J. Neurochem.* 82, 295–304.
- Mayer, M.L., and Westbrook, G.L. (1983). A voltage-clamp analysis of inward (anomalous) rectification in mouse spinal sensory ganglion neurones. *J. Physiol.* 340, 19–45.
- McLachlan, E.M., Jänig, W., Devor, M., and Michaelis, M. (1993). Peripheral nerve injury triggers noradrenergic sprouting within dorsal root ganglia. *Nature* 363, 543–546.
- Merkel, F. (1875). Tastzellen und tastkörperchen bei den hausthieren und beim menschen. *Archiv für Mikroskopische Anatomie* 11, 636–652.
- Mihara, M., Hashimoto, K., Ueda, K., and Kumakiri, M. (1979). The specialized junctions between Merkel cell and neurite: an electron microscopic study. *J. Invest. Dermatol.* 73, 325–334.
- Moehring, F., Cowie, A.M., Menzel, A.D., Weyer, A.D., Grzybowski, M., Arzua, T., Geurts, A.M., Palygin, O., and Stucky, C.L. (2018). Keratinocytes mediate innocuous and noxious touch via ATP-P2X4 signaling. *eLife* 7, eLife.31684.
- Molinoff, P.B., and Axelrod, J. (1971). Biochemistry of catecholamines. *Annu. Rev. Biochem.* 40, 465–500.
- Morita, T., McClain, S.P., Batia, L.M., Pellegrino, M., Wilson, S.R., Kienzler, M.A., Lyman, K., Olsen, A.S., Wong, J.F., Stucky, C.L., et al. (2015). HTR7 mediates serotonergic acute and chronic itch. *Neuron* 87, 124–138.
- Morrison, K.M., Miesegaes, G.R., Lumpkin, E.A., and Maricich, S.M. (2009). Mammalian Merkel cells are descended from the epidermal lineage. *Dev. Biol.* 336, 76–83.
- Nakamura, T., and Gold, G.H. (1987). A cyclic nucleotide-gated conductance in olfactory receptor cilia. *Nature* 325, 442–444.
- Nakamura, F., and Strittmatter, S.M. (1996). P2Y1 purinergic receptors in sensory neurons: contribution to touch-induced impulse generation. *Proc. Natl. Acad. Sci. USA* 93, 10465–10470.
- Niu, J., Vysochan, A., and Luo, W. (2014). Dual innervation of neonatal Merkel cells in mouse touch domes. *PLoS ONE* 9, e92027.
- Nunzi, M.G., Pisarek, A., and Mugnaini, E. (2004). Merkel cells, corpuscular nerve endings and free nerve endings in the mouse palatine mucosa express three subtypes of vesicular glutamate transporters. *J. Neurocytol.* 33, 359–376.
- Pang, Z., Sakamoto, T., Tiwari, V., Kim, Y.S., Yang, F., Dong, X., Güler, A.D., Guan, Y., and Caterina, M.J. (2015). Selective keratinocyte stimulation is sufficient to evoke nociception in mice. *Pain* 156, 656–665.
- Pennington, J.M., Millar, J., L Jones, C.P., Owesson, C.A., McLaughlin, D.P., and Stamford, J.A. (2004). Simultaneous real-time amperometric measurement of catecholamines and serotonin at carbon fibre ‘dident’ microelectrodes. *J. Neurosci. Methods* 140, 5–13.
- Press, D., Mutlu, S., and Güçlü, B. (2010). Evidence of fast serotonin transmission in frog slowly adapting type 1 responses. *Somatosens. Mot. Res.* 27, 174–185.
- R Development Core Team (2010). R: A language and environment for statistical computing (Vienna, Austria: R Foundation for Statistical Computing).
- Ramírez-Franco, J.J., Muñoz-Cuevas, F.J., Luján, R., and Jurado, S. (2016). Excitatory and inhibitory neurons in the hippocampus exhibit molecularly distinct large dense core vesicles. *Front. Cell. Neurosci.* 10, 202.
- Rausl, A., Nordlind, K., and Wahlgren, C.F. (2013). Pruritic and vascular responses induced by serotonin in patients with atopic dermatitis and in healthy controls. *Acta Derm. Venereol.* 93, 277–280.
- Reinisch, C.M., and Tschachler, E. (2005). The touch dome in human skin is supplied by different types of nerve fibers. *Ann. Neurol.* 58, 88–95.
- Rich, T.C., Fagan, K.A., Nakata, H., Schaack, J., Cooper, D.M., and Karpen, J.W. (2000). Cyclic nucleotide-gated channels colocalize with adenylyl cyclase in regions of restricted cAMP diffusion. *J. Gen. Physiol.* 116, 147–161.
- Rose, M.F., Ren, J., Ahmad, K.A., Chao, H.T., Klisch, T.J., Flora, A., Greer, J.J., and Zoghbi, H.Y. (2009). Math1 is essential for the development of hindbrain neurons critical for perinatal breathing. *Neuron* 64, 341–354.
- Rosenbaum, D.M., Rasmussen, S.G., and Kobilka, B.K. (2009). The structure and function of G-protein-coupled receptors. *Nature* 459, 356–363.
- Salzer, I., Gantumur, E., Yousuf, A., and Boehm, S. (2016). Control of sensory neuron excitability by serotonin involves 5HT2C receptors and Ca(2+)-activated chloride channels. *Neuropharmacology* 110 (Pt A), 277–286.
- Sato, J., and Perl, E.R. (1991). Adrenergic excitation of cutaneous pain receptors induced by peripheral nerve injury. *Science* 251, 1608–1610.
- Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* 9, 671–675.
- Shih, J.C., Chen, K., and Ridd, M.J. (1999). Monoamine oxidase: from genes to behavior. *Annu. Rev. Neurosci.* 22, 197–217.
- Shroyer, N.F., Helmrich, M.A., Wang, V.Y., Antalffy, B., Henning, S.J., and Zoghbi, H.Y. (2007). Intestine-specific ablation of mouse atonal homolog 1 (Math1) reveals a role in cellular homeostasis. *Gastroenterology* 132, 2478–2488.
- Stewart, W.C., and Castelli, W.P. (1996). Systemic side effects of topical beta-adrenergic blockers. *Clin. Cardiol.* 19, 691–697.
- Tachibana, T., and Nawa, T. (2002). Recent progress in studies on Merkel cell biology. *Anat. Sci. Int.* 77, 26–33.

- Tachibana, T., and Nawa, T. (2005). Immunohistochemical reactions of receptors to met-enkephalin, VIP, substance P, and CGRP located on Merkel cells in the rat sinus hair follicle. *Arch. Histol. Cytol.* *68*, 383–391.
- Torebjörk, E., Wahren, L., Wallin, G., Hallin, R., and Koltzenburg, M. (1995). Noradrenaline-evoked pain in neuralgia. *Pain* *63*, 11–20.
- Toyoshima, K., and Shimamura, A. (1991). Uranaffin reaction of Merkel corpuscles in the lingual mucosa of the finch, *Lonchula striata* var. *domestica*. *J. Anat.* *179*, 197–201.
- Urtikova, N., Berson, N., Van Steenwinckel, J., Doly, S., Truchetto, J., Maroteaux, L., Pohl, M., and Conrath, M. (2012). Antinociceptive effect of peripheral serotonin 5-HT_{2B} receptor activation on neuropathic pain. *Pain* *153*, 1320–1331.
- Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., Hjerling-Leffler, J., Haeggström, J., Kharchenko, O., Kharchenko, P.V., et al. (2015). Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nat. Neurosci.* *18*, 145–153.
- Van Buskirk, E.M. (1980). Adverse reactions from timolol administration. *Ophthalmology* *87*, 447–450.
- Van Keymeulen, A., Mascré, G., Youseff, K.K., Harel, I., Michaux, C., De Geest, N., Szpalski, C., Achouri, Y., Bloch, W., Hassan, B.A., and Blanpain, C. (2009). Epidermal progenitors give rise to Merkel cells during embryonic development and adult homeostasis. *J. Cell Biol.* *187*, 91–100.
- Vielkind, U., Sebzda, M.K., Gibson, I.R., and Hardy, M.H. (1995). Dynamics of Merkel cell patterns in developing hair follicles in the dorsal skin of mice, demonstrated by a monoclonal antibody to mouse keratin 8. *Acta Anat. (Basel)* *152*, 93–109.
- Weihe, E., Hartschuh, W., Schäfer, M.K., Romeo, H., and Eiden, L.E. (1998). Cutaneous Merkel cells of the rat contain both dynorphin A and vesicular monoamine transporter type 1 (VMAT1) immunoreactivity. *Can. J. Physiol. Pharmacol.* *76*, 334–339.
- Wellnitz, S.A., Lesniak, D.R., Gerling, G.J., and Lumpkin, E.A. (2010). The regularity of sustained firing reveals two populations of slowly adapting touch receptors in mouse hairy skin. *J. Neurophysiol.* *103*, 3378–3388.
- Westlund, K.N., and Coulter, J.D. (1980). Descending projections of the locus coeruleus and subcoeruleus/medial parabrachial nuclei in monkey: axonal transport studies and dopamine-beta-hydroxylase immunocytochemistry. *Brain Res.* *2*, 235–264.
- Wilson, S.R., Thé, L., Batia, L.M., Beattie, K., Katibah, G.E., McClain, S.P., Pellegrino, M., Estandian, D.M., and Bautista, D.M. (2013). The epithelial cell-derived atopic dermatitis cytokine TSLP activates neurons to induce itch. *Cell* *155*, 285–295.
- Woo, S.H., Ranade, S., Weyer, A.D., Dubin, A.E., Baba, Y., Qiu, Z., Petrus, M., Miyamoto, T., Reddy, K., Lumpkin, E.A., et al. (2014). Piezo2 is required for Merkel-cell mechanotransduction. *Nature* *509*, 622–626.
- Woodbury, C.J., and Koerber, H.R. (2007). Central and peripheral anatomy of slowly adapting type I low-threshold mechanoreceptors innervating trunk skin of neonatal mice. *J. Comp. Neurol.* *505*, 547–561.
- Yamamoto, M., Wada, N., Kitabatake, Y., Watanabe, D., Anzai, M., Yokoyama, M., Teranishi, Y., and Nakanishi, S. (2003). Reversible suppression of glutamatergic neurotransmission of cerebellar granule cells in vivo by genetically manipulated expression of tetanus neurotoxin light chain. *J. Neurosci.* *23*, 6759–6767.
- Yau, K.W., and Baylor, D.A. (1989). Cyclic GMP-activated conductance of retinal photoreceptor cells. *Annu. Rev. Neurosci.* *12*, 289–327.
- Yu, C.R., Power, J., Barnea, G., O'Donnell, S., Brown, H.E., Osborne, J., Axel, R., and Gogos, J.A. (2004). Spontaneous neural activity is required for the establishment and maintenance of the olfactory sensory map. *Neuron* *42*, 553–566.
- Zeitz, K.P., Guy, N., Malmberg, A.B., Dirajlal, S., Martin, W.J., Sun, L., Bonhaus, D.W., Stucky, C.L., Julius, D., and Basbaum, A.I. (2002). The 5-HT₃ subtype of serotonin receptor contributes to nociceptive processing via a novel subset of myelinated and unmyelinated nociceptors. *J. Neurosci.* *22*, 1010–1019.
- Zhang, Y., Narayan, S., Geiman, E., Lanuza, G.M., Velasquez, T., Shanks, B., Akay, T., Dyck, J., Pearson, K., Gosgnach, S., et al. (2008). V3 spinal neurons establish a robust and balanced locomotor rhythm during walking. *Neuron* *60*, 84–96.
- Zhou, F.W., Dong, H.W., and Ennis, M. (2016). Activation of β -noradrenergic receptors enhances rhythmic bursting in mouse olfactory bulb external tufted cells. *J. Neurophysiol.* *116*, 2604–2614.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rat monoclonal anti-KRT8	Developmental Studies Hybridoma Bank	Cat# TROMA-I; RRID: AB_531826
Chicken polyclonal anti-NFH	Abcam	Cat# ab4680; RRID: AB_304560
Chicken polyclonal anti-GFP	Abcam	Cat# ab13970; RRID: AB_300798
Rabbit polyclonal anti-NFH	Abcam	Cat# ab8135; RRID: AB_306298
Rabbit polyclonal anti-VAMP2	Abcam	Cat# ab3347; RRID: AB_2212462
Rabbit polyclonal anti- β 2-AR	Santa Cruz Biotechnology	Cat# sc-569; RRID: AB_630926
Rabbit polyclonal anti-TH	Millipore	Cat# AB152; RRID: AB_390204
Rabbit polyclonal anti-DBH	Immunostar	Cat# 22806; RRID: AB_572229
Rabbit polyclonal anti-DsRed	Clontech Laboratories	Cat# 632496; RRID: AB_10013483
Goat anti-Rat polyclonal Alexa 488	Thermo Fisher Scientific	Cat# A-11006; RRID: AB_141373
Goat anti-Rat polyclonal Alexa-594	Thermo Fisher Scientific	Cat# A-11007; RRID: AB_10561522
Goat anti-Rat polyclonal Alexa-647	Thermo Fisher Scientific	Cat# A-21247; RRID: AB_141778
Goat anti-Chicken polyclonal Alexa 488	Thermo Fisher Scientific	Cat# A-11039; RRID: AB_142924
Goat anti-Chicken polyclonal Alexa-594	Thermo Fisher Scientific	Cat# A-11042; RRID: AB_142803
Goat anti-Chicken polyclonal Alexa-647	Thermo Fisher Scientific	Cat# A-21449; RRID: AB_2535866
Goat anti-Rabbit polyclonal Alexa 488	Thermo Fisher Scientific	Cat# A-11008; RRID: AB_143165
Goat anti-Rabbit polyclonal Alexa-594	Thermo Fisher Scientific	Cat# A-11037; RRID: AB_2534095
Goat anti-Rabbit polyclonal Alexa-647	Thermo Fisher Scientific	Cat# A-21450; RRID: AB_141882
Chemicals, Peptides, and Recombinant Proteins		
Dispase	Thermo Fisher Scientific	Cat# CB-40235
O.C.T. Compound	Tissue-Tek	Cat# 4583
Triton X-100	Sigma-Aldrich	Cat# T9284
Fluoromount-G	Thermo Fisher Scientific	Cat# 00-4958-02
FM1-43	Biotium	Cat# 70022
Norepinephrine	Tocris	Cat# 5169
Serotonin Hydrochloride	Tocris	Cat# 3547
Dopamine Hydrochloride	Sigma-Aldrich	Cat# H8502
ICI 118,551 Hydrochloride	Tocris	Cat# 0821
Fura-2, AM	Thermo Fisher Scientific	Cat# F1221
Pluronic F-127	Thermo Fisher Scientific	Cat# P3000MP
ω -Conotoxin MVIIC	Sigma-Aldrich	Cat# C4188
Nimodipine	Tocris	Cat# 0600
FFN206	Tocris	Cat# 5043
Reserpine	Tocris	Cat# 2742
Critical Commercial Assays		
RNeasy Mini Kit	QIAGEN	Cat# 74104
Ovation RNA-Seq System V2	Nugen	Cat# 7102-32
Ovation Ultralow System V2	Nugen	Cat# 0344-32
RNAscope Fluorescent Multiplex kit	Advanced Cell Diagnostics	Cat# 320850
Deposited Data		
RNA-seq Data	This paper	GEO: GSE114336
Mendeley Dataset	This paper	https://doi.org/10.17632/pr87ymnmjm.2

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains		
Mouse: <i>Atoh1</i> ^{GFP} : B6.129S-Atoh1 ^{tm4.1Hzo} /J	Rose et al., 2009	RRID: MGI:4418333
Mouse: <i>Th</i> ^{GFP} : Tg(Th-EGFP)21-31Koba	Matsushita et al., 2002	RRID: IMSR_RBRC02095
Mouse: C57BL/6J	The Jackson Laboratory	RRID: IMSR_JAX:000664
Mouse: <i>K14</i> ^{Cre} : Tg(KRT14-cre)1Amc/J	The Jackson Laboratory	RRID: IMSR_JAX:004782
Mouse: <i>Atoh1</i> ^{LacZ} : B6.129S7-Atoh1 ^{tm2Hzo} /J	The Jackson Laboratory	RRID: IMSR_JAX:005970
Mouse: <i>Atoh1</i> ^{fllox} : B6.129S7-Atoh1 ^{tm3Hzo} /J	The Jackson Laboratory	RRID: IMSR_JAX:008681
Mouse: 5HT _{3A} -EGFP: Tg(Htr3a-EGFP)	Gong et al., 2003	RRID: MMRRC_000273-UNC
Mouse: <i>Atoh1</i> ^{nGFP} : Tg(Atoh1-GFP)1Jejo	Lumpkin et al., 2003	MGI:3703598
Mouse: <i>R26</i> ^{TeNT} : Gt(ROSA)26Sor ^{tm2(GFP/tetX)Gld}	Zhang et al., 2008	MGI:3839913
Mouse: <i>Wnt1</i> ^{Cre} : 129S4.Cg-Tg(Wnt1-cre)2Sor/J	The Jackson Laboratory	RRID: IMSR_JAX:022137
Mouse: <i>TrkC</i> ^{tdTomato} : Ntrk3 ^{tm2.1Ddg}	Bai et al., 2015	RRID: MGI:5707229
Mouse: <i>Adrb2</i> ^{fllox} : <i>Adrb2</i> ^{tm1Kry}	Hinoi et al., 2008	MGI:3837346
Oligonucleotides		
<i>Adrb2</i> probe channel 3	Advanced Cell Diagnostics	Cat# 449771-C3
Software and Algorithms		
SAMtools	Li et al., 2009	http://samtools.sourceforge.net
R v3.4.3	R Development Core Team, 2010	http://www.R-project.org
DESeq2	Love et al., 2014	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
ImageJ	Schneider et al., 2012	https://imagej.nih.gov/ij/index.html
MATLAB 2017b	MathWorks	http://www.mathworks.com
Custom Spike Sorting Software	Hoffman and Lumpkin, 2018	https://github.com/buh2003/SpikeSortingPCA_DBSCAN
Metafluor v7.6.3	Molecular Devices	https://www.moleculardevices.com
Prism v7.0b	GraphPad	http://www.graphpad.com

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to Dr. Ellen A. Lumpkin (ellen.a.lumpkin@gmail.com).

EXPERIMENTAL MODEL AND SUBJECT DETAILS**Animals**

Animal use was conducted according to guidelines from the National Institutes of Health's Guide for the Care and Use of Laboratory Animals and was approved by the Institutional Animal Care and Use Committee of Columbia University Medical Center. Mice were maintained on a 12 h light/dark cycle, and food and water was provided *ad libitum*. All mice were healthy, and none of the experimental mice was immune compromised. Unless otherwise noted, mice of both sexes were used and mice were randomly allocated to experimental groups. The following strains were used in this study: *Atoh1*^{GFP} (Rose et al., 2009), *Th*^{GFP} (Matsushita et al., 2002) which express GFP under the control of the *Th* promoter, wild-type (WT) C57BL/6J, *K14*^{Cre} (Dassule et al., 2000), *Atoh1*^{LacZ} (Ben-Arie et al., 2000), *Atoh1*^{fllox} (Shroyer et al., 2007), 5HT_{3A}-EGFP (Gong et al., 2003), *Atoh1*^{nGFP} (Lumpkin et al., 2003), *R26*^{TeNT} (Zhang et al., 2008), *Wnt1*^{Cre} (Lewis et al., 2013), *TrkC*^{tdTomato} (Bai et al., 2015), and *Adrb2*^{fllox} (Hinoi et al., 2008). *TeNT* mice were generated by crossing hemizygous *K14*^{Cre} male with homozygous *R26*^{TeNT} female mice. *Atoh1*^{CKO} mice were generated as described previously (Morrison et al., 2009). Mice of the *K14*^{Cre}; *Atoh1*^{LacZ/fllox} genotype lacked expression of *Atoh1* in *K14*-expressing cells and were designated as *Atoh1*^{CKO} mice. Genotypes that lacked either the Cre and/or LacZ alleles were designated as controls. *Adrb2*^{CKO} mice were generated by first crossing hemizygous *Wnt1*^{Cre} male with homozygous *Adrb2*^{fllox} mice. Male progeny hemizygous for *Wnt1*^{Cre} and heterozygous for *Adrb2*^{fllox} (*Wnt1*^{Cre}; *Adrb2*^{fllox/+}) were crossed with homozygous *Adrb2*^{fllox} female mice. Mice of the *Wnt1*^{Cre}; *Adrb2*^{fllox/fllox} genotype were designated as experimental, and mice with the *Adrb2*^{fllox/fllox} that lacked Cre were designated as controls. All mice were bred as recommended by The Jackson Laboratory. Genotyping was performed through a commercial service (Transnetyx).

METHOD DETAILS

RNA Sequencing and Analysis

To isolate Merkel cells and keratinocytes for RNA-seq, dorsal skin from 7–8-week old female mice was removed and placed in a plastic dish, epidermis-side down. Skin was scraped with a scalpel to remove fat and muscle. Skin was then floated epidermis-side up in 0.25% trypsin for 2 h at 37°C. Epidermis was manually separated from the dermis, broken into smaller pieces, and recovered in culture media (CnT02, Chemicon) for 30 min with a stir bar. A 70- μ m cell strainer was used to collect single cells. Cells were incubated with the Cd49f-PE antibody (BD PharMingen) for 30 min on ice. GFP+/PE- Merkel cells and GFP-/PE+ keratinocytes were purified from the epidermal-cell suspension with FACS, directly into Trizol (ThermoFisher). Total RNA was isolated using commercially available reagents (QIAGEN RNeasy kit) and DNase-treated according to manufacturer's instructions to remove contaminating genomic DNA. First- and second-strand cDNA synthesis, and cDNA amplification was performed with the Ovation RNA-Seq System V2 (NuGEN). The cDNA library was prepared with the Ovation Ultralow System V2 (NuGEN). Sequencing was conducted on an Illumina HiSeq1000. Reads were aligned to the annotated mouse reference genome (mm10) with SAMtools (Li et al., 2009). Differential expression analysis was performed with DESeq2 (Love et al., 2014). The core presynaptic gene list (Table S2) was generated by combining published presynaptic proteomic libraries (Abul-Husn et al., 2009; Boyken et al., 2013) with the MGI (Blake et al., 2017) and Ensembl (Aken et al., 2017) presynapse (GO:0098793) gene ontology annotations (combined, 542 genes). Genes were manually annotated according to protein function using Gene Ontology annotations.

Semi-intact Epidermal Peel Preparation

Live-cell imaging was performed in a semi-intact epidermal peel preparation. This preparation enables direct pharmacological and physiological access to Merkel cells *in situ*. Back skin from P20–24 *Atoh1^{hGFP}* or *Atoh1^{GFP}* mice was clipped with electric clippers, depilated (Surgi-cream), and dissected onto a small dish. Skin specimens (1 cm²) were applied to glass coverslips epidermis-side down, in order to form a stable adhesion between the coverslip and epidermis. Samples were incubated in dispase (25 U/ml, Fisher Scientific) for 1 h at room temperature, on an orbital shaker. Using forceps, dermal and subcutaneous tissue was gently peeled from the epidermis, leaving only epidermal cells on the coverslip.

Immunohistochemistry

For cryosections, mouse skin was clipped with electric clippers, depilated (Surgi-cream) and dissected from the back (7–10-weeks of age). Tissue was fixed in 4% paraformaldehyde (PFA) for 30 min, cryoprotected in 30% sucrose overnight, frozen in OCT (Tissue-Tek), and sectioned at a thickness of 16–20 μ m. For epidermal peel preparations, samples were fixed in 4% PFA for 30 min. Cryosectioned skin or epidermal peels were labeled at 4°C overnight with the following primary antibodies: rat anti-K8 (TROMA1, Developmental Studies Hybridoma Bank, 1:100), chicken anti-NFH (Abcam: ab4680, 1:2,000), rabbit anti-VAMP2 (Abcam: ab3347, 1:500), rabbit anti- β 2-AR (H-20, Santa Cruz: sc-569, 1:250), rabbit anti-TH (Millipore: AB152, 1:500), and rabbit anti-DBH (Immunostar: 22806, 1:2000). Secondary goat AlexaFluor-conjugated antibodies (Thermo Fisher Scientific) directed against rat (Alexafluor 488, A-11006; Alexafluor 594, A-11007), chicken (Alexafluor 488, A-11039; Alexafluor 594, A-11042), or rabbit (Alexafluor 488, A-11008; Alexafluor 647, A-21450) IgG were used for 1 h at room temperature (1:1000). Samples were mounted with Fluoromount-G with DAPI (Thermo Fisher Scientific).

For whole-mount immunostaining with tissue clearing, skin was depilated, tape-stripped to remove the stratum corneum and dissected from the proximal hind limb of mice (7–10 weeks of age). For whisker whole-mount immunostaining, whiskers were isolated from whisker pads of mice (7–10 weeks of age), and whisker capsules were dissected to expose the ring sinus and external root sheath. Ring sinus was removed to expose the glassy membrane and Merkel cells within. Whole-mount immunohistochemistry was performed as previously described (Lesniak et al., 2014). Briefly, tissue was fixed in 4% PFA overnight, washed in 0.03% triton-X PBS (PBST) and incubated in primary antibody for 72–96 h at 4°C. Primary antibodies used were: rat anti-K8 (TROMA1, Developmental Studies Hybridoma Bank, 1:100), chicken anti-NFH (Abcam: ab4680, 1:500), rabbit anti-NFH (Abcam; ab8135, 1:500), and chicken anti-GFP (Abcam: ab13970, 1:500). After 5–10 h of washes in PBST, samples were incubated for 48 h at 4°C in secondary antibodies: goat AlexaFluor-conjugated antibodies (Thermo Fisher Scientific) directed against rat (Alexafluor 488, A-11006; Alexafluor 647, A-21247), chicken (Alexafluor 488, A-11039; Alexafluor 594, A-11042) or rabbit (Alexafluor 594, A-11037) IgG. After staining, tissue was dehydrated progressively in 25%, 50%, 75% and 100% methanol in PBST (1 h each) and cleared using a 2:1 benzyl benzoate/benzyl alcohol solution.

Specimens were imaged in three dimensions (0.5–1 μ m axial step sizes) on a Zeiss Exciter confocal microscope (LSM 5) equipped with 20X, 0.8 NA or 40X, 1.3 NA objective lenses.

In Situ Hybridization

DRG sections harvested from *TrkC^{tdTomato}* mice (4 weeks of age) were cut at 25- μ m thickness and processed for high-sensitivity RNA *in situ* detection using an RNAscope Fluorescent Detection Kit according to manufacturer's instructions (Advanced Cell Diagnostics, Hayward, California, USA). The following modifications were made to the protocol: after harvesting, DRG were fixed in 4% PFA for 15 min and then incubated in 30% sucrose for 2 h at 4°C. DRG were embedded in OCT (Sakura) and stored at –80°C until sectioned. Additional fixation steps were omitted from hybridization protocol. Hybridization was performed on DRG sections using the *Adrb2*

probe (449771-C3, mouse), followed by incubation at 4°C overnight with rabbit anti-dsRed (Clontech: 632496, 1:3000) and chicken anti-NFH (Abcam: ab4680, 1:5000) primary antibodies. Sections were then incubated with goat AlexaFluor-conjugated antibodies directed against rabbit (Thermo Fisher Scientific, Alexafluor 594, A-11037) and chicken (Alexafluor 647, A-21449) at room temperature for 1 h at a concentration of 1:1000. Samples were mounted with Fluoromount-G (Fisher Scientific). Specimens were imaged in three dimensions (1- μ m axial steps) on a Nikon Ti Eclipse for scanning confocal microscopy equipped with a 40X, 1.3 NA objective lens. Images were analyzed using ImageJ software (Schneider et al., 2012). Quantification was performed on unprocessed axial stacks, following thresholding.

High-Performance Liquid Chromatography (HPLC)

Whisker pads were dissected from *Atoh1*^{CKO} and littermate control animals (7–10 weeks of age) and placed in a dissection dish with PBS. Samples were chilled on ice to prevent monoamine oxidation. Follicles were exposed in whisker pads for further micro-dissection. To reduce vascular contributions, follicles were trimmed at the junction between the ring and cavernous sinuses. Once trimmed, the ring sinus was removed to expose the glassy membrane and the Merkel cells within the external root sheath. Trimmed follicles were then separated from whisker pads by transection at the inner conical body. Micro-dissected follicles were immediately placed in a 0.1 M perchloric acid solution on ice. Four follicles were pooled per animal. Pooled follicles were homogenized with a hand held sonicator and centrifuged (15,000 rpm) at 4°C. Supernatant was then stored at –80°C or analyzed immediately. HPLC was performed as described previously (Feigin et al., 2001; Larsen et al., 2002). Briefly, HPLC coupled with electrochemical detection was performed on an ESA Coulochem II detector equipped with a model 5011 analytical cell (ESA) set at an applied potential of 400 mV and a Velosep RP-18 column (Applied Biosystems). The mobile phase contained 45 mM NaH₂PO₄, 0.2 mM EDTA, 1.2 mM heptanesulfonic acid and 5% methanol, adjusted to pH 3.2 with phosphoric acid. Total elution time was 30 min. Molar amounts of metabolites were calculated from areas under HPLC peaks using calibration curves and normalized to total sample weight.

FM1-43 Injections

FM1-43 (Biotium; #70020) was used to visualize SAI receptive fields (touch domes) for electrophysiological recordings. FM1-43 was diluted at 1.5 mM in sterile PBS and injected subcutaneously (70 μ l per mouse). Tissue was dissected for *ex vivo* skin–nerve electrophysiology 12–14 h after injection.

Ex Vivo Skin-Nerve Electrophysiology

Touch-evoked responses from cutaneous afferents were recorded after dissecting the hindlimb skin and saphenous nerve from 7–13 week old mice, according to published methods (Wellnitz et al., 2010). For identifying receptive fields, the skin was placed epidermis-side-up in a custom chamber and perfused with carbogen-buffered synthetic interstitial fluid (SIF) kept at 32°C with a temperature controller (model TC-344B, Warner Instruments). The nerve was immersed in mineral oil in a recording chamber, teased apart with fine forceps, and small bundles were placed onto a silver recording electrode connected with a reference electrode to a differential amplifier (model 1800, A-M Systems). Conduction velocity was measured by electrically stimulating receptive fields (Model 2100 isolated pulse stimulator, A-M Systems). Signals were band-passed filtered at 0.3–5 kHz, sampled at 20 kHz using a PowerLab 8/35 board (AD Instruments) and recorded using LabChart software (AD Instruments). SAI receptive fields (touch domes) labeled with FM1-43 were visualized using a fluorescence stereomicroscope equipped with a long-pass GFP filter set.

Spike sorting and data analysis was performed offline with custom software in MATLAB (Hoffman and Lumpkin, 2018). Semi-supervised spike sorting was performed with principal component analysis (PCA) and density-based spatial clustering of applications with noise (DBSCAN). Sorted spikes were then analyzed for metrics of spike timing, firing rate, and adaptation properties.

For these studies, we focused on Merkel-cell afferents in touch domes, as described by Iggo and Muir (Iggo and Muir, 1969). The afferents generally have no spontaneous firing, respond selectively to pressure applied directly to a touch dome, and are particularly sensitive to moving stimuli but are insensitive to hair tugging and skin stretch (Iggo and Muir, 1969). To identify responses from these afferents in mutant and control genotypes, we used a mechanical search paradigm with a fine glass probe. Afferents were classified as ‘Merkel-cell afferents’ according to the following criteria: (1) A β conduction velocity (≥ 9 m s⁻¹), (2) punctate receptive fields restricted to one or more fluorescently labeled touch domes, (3) insensitive to pressure applied to skin areas adjacent to touch domes, (4) insensitive to hair tugging but responsive when the hair is bent to compress the touch dome. Touch-sensitive afferents that did not meet these criteria were not analyzed further. Fibers were classified as SA if spikes were observed during the last 1 s of the hold phase of stimulation in $\geq 75\%$ of stimulus presentations, otherwise they were classified as IA. Recordings and analyses of *K14*^{Cre}; *R26*^{TeNT}, *Wnt1*^{Cre}; *Adrb2*^{fl/fl}, and their controls were performed blind to genotype (Table S3).

Afferents were classified as A-fiber RA-LTMRs according to the following criteria: (1) A-fiber conduction velocity (> 1 m s⁻¹), (2) rapid adaptation to mechanical indentation, (3) sensitive to low-force stimulation with von Frey monofilaments (< 0.4 mN).

Mechanical responses were elicited with von Frey monofilaments and a custom-built mechanical stimulator. The automated mechanical stimulator applied stimuli with an indenter (tip diameter, 1.6 mm), and stimuli were commanded using a model XPS motion controller and driver system (Newport) connected to a PC computer. Movement of the indenter was controlled with custom software and measured with a laser distance-measuring device (OptiNCDT 1402, Micro-Epsilon). Touch stimuli consisted of ramp and 5 s hold indentations. First-order approximation of approach speed was 3.2 mm s⁻¹. The mechanical stimulator tip was

held ~600 μm from the surface of the skin. Mechanical displacements ranged from the point of skin contact to 450 μm of skin indentation. For non-pharmacological experiments, the period between successive displacements was 60 s. For pharmacological experiments, the period between successive displacements was 120 s.

For pharmacology, receptive fields, conduction velocity and von Frey thresholds were first identified with the epidermis-side facing up. After identification of a Merkel-cell afferent based on FM1-43 fluorescence and the SAI response pattern, the skin was inverted such that the dermis side faced up for perfusion. Receptive fields were then isolated with a custom-built glass perfusion ring. For experiments performed in the absence of mechanical stimulation, drugs were directly applied to the isolated receptive fields (Norepinephrine, Tocris, 5169; Serotonin, Tocris, 3547; Dopamine, Sigma Aldrich, H8502) and evoked responses were recorded in gap-free mode. For experiments performed with simultaneous pharmacological application and mechanical stimulation, both drugs and mechanical indentation were applied to receptive fields within the perfusion ring (ICI 118,551, Tocris, 0821). To accommodate for repeated mechanical stimulations over long periods of time, we reduced the magnitude of mechanical stimuli for pharmacological experiments. Thus, recorded firing rates from pharmacological experiments were slightly lower than in non-pharmacological experiments (compare Figures 3D–3F to Figures 5B–5F).

Norepinephrine-response data were fit with the following four-parameter logistic equation:

$$y = \frac{a - b}{1 + 10^{((\log_{10} c) - x) \times d)} + b}$$

$a = \text{maximum}; b = \text{minimum}; c = EC_{50}; d = \text{Hill's slope}$

Calcium Imaging

Semi-intact epidermal peel preparations were isolated from *Atoh1^{nGFP}* or *Atoh1^{GFP}* mice. Merkel cells were loaded for 30 min at 37°C with 5 μM fura-2 acetoxymethyl ester (Thermo Fisher Scientific) and 1 μM pluronic acid (Pluronic F-127, Thermo Fisher Scientific) in a modified Ringer's solution (in mM): 140 NaCl, 5 KCl, 10 HEPES (pH 7.4), 10 D-Glucose, 2 MgCl_2 , and 2 CaCl_2 (osmolality: 290 mmol kg^{-1}). Cells were given 30 min in Ringers solution to digest the ester bonds before imaging. Merkel cells were depolarized with mechanical stimulation or with high potassium extracellular solution (in mM): 5 NaCl, 140 KCl, 10 HEPES (pH 7.4), 10 D-Glucose, 2 MgCl_2 , and 2 CaCl_2 (osmolality: 290 mmol kg^{-1}). Mechanical stimulation was delivered with a glass probe (tip diameter: ~0.5 μm) driven with a stepper motor (model MP-200, Sutter Instruments). The glass probe was positioned at an angle of 40° to the coverslip. To ensure that mechanosensitive channels were not activated at rest, displacements began from an offset position located 2–3 μm away from Merkel cells (Drew et al., 2002). For experiments with voltage-gated calcium channel blockers (nimodipine, Tocris, 0600; ω -Conotoxin MVIIC, Sigma Aldrich, C4188), Merkel cells were incubated with blockers for 30 min before imaging. Data were acquired with Metafluo software (version 7.6.3, Molecular Devices), and analyzed with ImageJ. Briefly, xy shifts were corrected in 340-nm and 380-nm images using a rigid body transformation with the “StackReg” plugin (Plugins/Registration/StackReg/RigidBody). Images of the fluorescence ratio at 340 nm and 380 nm excitation ($F_{340/380}$) were created with the “Image Calculator” (Image/Image Calculator). Regions of interest (ROIs) were drawn based on identified Merkel cells. Mean ROI fluorescence intensity over time was quantified with the “Time Series Analyzer V3” (Plugins/Time Series Analyzer V3). $\Delta F/F$ values were calculated as follows: $(F - F_i)/F_i$, where F is the $F_{340/380}$ at each frame and F_i is the average $F_{340/380}$ for the 30 s prior to stimulation. Samples were imaged with a 20x, 0.95 NA objective lens.

FFN Imaging

Semi-intact epidermal peel preparations were isolated from *Atoh1^{nGFP}* or *Atoh1^{GFP}* mice. Merkel cells were loaded for 30 min at 37°C with 1 μM FFN206 (Tocris) in a modified Ringer's solution (in mM): 140 NaCl, 5 KCl, 10 HEPES (pH 7.4), 10 D-Glucose, 2 MgCl_2 , and 2 CaCl_2 (osmolality: 290 mmol kg^{-1}). Merkel cells were depolarized with mechanical stimulation or with high potassium extracellular solution (in mM): 5 NaCl, 140 KCl, 10 HEPES (pH 7.4), 10 D-Glucose, 2 MgCl_2 , and 2 CaCl_2 (osmolality: 290 mmol kg^{-1}). Mechanical stimulation was delivered with a glass probe (tip diameter: ~0.5 μm) driven by a piezoelectric actuator (model PA8/12, Piezosystem Jena; power supply ENV40 C, Piezosystem Jena). The glass probe was positioned at an angle of 40° to the coverslip. Displacements were triggered by a pClamp-controlled command voltage passed to the actuator driver through a low-pass filter (f_{cutoff} : 500 Hz; model LPF-100A, Warner Instruments). Displacement magnitudes were visually calibrated daily. To ensure that mechanosensitive channels were not activated at rest, displacements began from an offset position located 2–3 μm away from Merkel cells (Drew et al., 2002). For experiments with voltage-gated calcium channel blockers (nimodipine, Tocris, 0600; ω -Conotoxin MVIIC, Sigma Aldrich, C4188), Merkel cells were incubated with blockers for 30 min before imaging. Data were acquired with Metafluo software (version 7.6.3, Molecular Devices).

ImageJ and MATLAB were used to analyze FFN206 destaining experiments. First, xy shifts were corrected in ImageJ using a rigid body transformation with the “StackReg” plugin (Plugins/Registration/StackReg/RigidBody). Next, ROIs were drawn based on FFN206 puncta within GFP-positive Merkel cells. Mean ROI fluorescence intensity over time was quantified with the “Time Series Analyzer V3” (Plugins/Time Series Analyzer V3). All further analyses were performed in MATLAB. To correct for bleaching,

the baseline intensity values for each ROI were individually fit with a monoexponential decay function. To obtain bleaching-corrected values, fluorescence values (F) were subtracted from the monoexponential fit (F_{exp}) for that ROI ($F_{corr} = 1 + F - F_{exp}$). Baseline normalized values (F/F_i) were calculated by dividing bleaching-corrected fluorescence values (F) for each time point by the mean of the first 30 s of each ROI (F_i). Puncta were classified as “destaining” if they exhibited at least 2% decrease in (F/F_i) 240 s after high K^+ stimulation or 40 s after mechanical stimulation. Samples were imaged with a 20x, 0.95 NA objective lens (Olympus).

QUANTIFICATION AND STATISTICAL ANALYSIS

Analysis was performed in R (R-Project; [R Development Core Team, 2010](#)), MATLAB (MathWorks), and Prism (Graphpad). Detailed statistical parameters are described in figure legends. Data are represented as mean \pm SEM and n represents the number of cells or independent experiments. For parametric data with three or more groups, one-way or two-way ANOVAs were followed by post hoc analyses for between-group comparisons. Unpaired, non-parametric data were analyzed using the Kruskal-Wallis test with Dunn’s post hoc comparison. Paired data were compared with a matched two-way ANOVA with Sidak’s post hoc. Student’s two-tailed t test was used to compare means of two normally distributed groups. Categorical data were compared using the two-tailed Fisher’s exact test. The normality of population data was assessed using the Kolmogorov-Smirnov test with Dallal-Wilkinson-Lilliefors p values, with $p < 0.05$ indicating non-normality. Data were considered significant if $p < 0.05$. Statistical details of experiments can be found in figure legends.

DATA AND SOFTWARE AVAILABILITY

The RNA-seq data are accessible at Gene Expression Omnibus (GEO: GSE114336). The source data for this paper are accessible at <https://doi.org/10.17632/pr87ymnmjm.2>. Spike sorting and *ex vivo* data analysis software is freely available at http://www.github.com/buh2003/spikesortingPCA_DBSCAN. All other data are available from the authors upon request.